



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# HSP 56 (h3): 293T Lysate: sc-158624

## BACKGROUND

HSP 56 (also designated FKBP4, FK506 binding protein 4, HBI, p52, FKBP52, FKBP59 and PPlase) is a *cis-trans* prolyl isomerase belonging to the immunophilin protein family. The human HSP 56 gene (FKBP4) has multiple polyadenylation sites and the HSP 56 protein can undergo phosphorylation. HSP 56 influences immunoregulatory gene expression in lymphocytes, protein folding and trafficking. It can serve as a co-chaperone for steroid hormone nuclear receptors to govern appropriate hormone action in target tissues. The protein can associate with phytanoyl-CoA  $\alpha$ -hydroxylase (PHYH) and with HSP 90 through a series of tetratricopeptide repeat (TPR) domains. HSP 56 is a TRPC ion channel accessory protein that modulates channel activation following receptor stimulation.

## REFERENCES

1. Sumanasekera, W.K., et al. 2003. Heat shock protein 90 (HSP 90) acts as a repressor of peroxisome proliferator-activated receptor  $\alpha$  (PPAR $\alpha$ ) and PPAR $\beta$  activity. *Biochemistry* 42: 10726-10735.
2. Galigniana, M.D., et al. 2004. HSP 90-binding immunophilins link p53 to Dynein during p53 transport to the nucleus. *J. Biol. Chem.* 279: 22483-22489.
3. Sanokawa-Akakura, R., et al. 2004. A novel role for the immunophilin FKBP52 in copper transport. *J. Biol. Chem.* 279: 27845-27848.
4. Sinkins, W.G., et al. 2004. Association of immunophilins with mammalian TRPC channels. *J. Biol. Chem.* 279: 34521-34529.
5. Davies, T.H., et al. 2005. FKBP52. *Int. J. Biochem. Cell Biol.* 37: 42-47.
6. Wochnik, G.M., et al. 2005. FK506-binding proteins 51 and 52 differentially regulate Dynein interaction and nuclear translocation of the glucocorticoid receptor in mammalian cells. *J. Biol. Chem.* 280: 4609-4616.
7. Prince, T., et al. 2005. Evidence for chaperone heterocomplexes containing both HSP 90 and VCP. *Biochem. Biophys. Res. Commun.* 331: 1331-1337.
8. Tranguch, S., et al. 2005. Co-chaperone immunophilin FKBP52 is critical to uterine receptivity for embryo implantation. *Proc. Natl. Acad. Sci. USA* 102: 14326-14331.
9. Kim, S.H., et al. 2005. Mitogenic estrogen metabolites alter the expression of 17 $\beta$ -estradiol-regulated proteins including heat shock proteins in human MCF7 breast cancer cells. *Mol. Cells* 20: 378-384.

## CHROMOSOMAL LOCATION

Genetic locus: FKBP4 (human) mapping to 12p13.33.

## PRODUCT

HSP 56 (h3): 293T Lysate represents a lysate of human HSP 56 transfected 293T cells and is provided as 100  $\mu$ g protein in 200  $\mu$ l SDS-PAGE buffer.

## STORAGE

Store at -20 $^{\circ}$  C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

## APPLICATIONS

HSP 56 (h3): 293T Lysate is suitable as a Western Blotting positive control for human reactive HSP 56 antibodies. Recommended use: 10-20  $\mu$ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.