



**SZABO  
SCANDIC**

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](http://linkedin.com/company/szaboscandic)



# pancpin (h2): 293 Lysate: sc-158810



## BACKGROUND

Pancpin, also known as myoepithelium-derived serine protease inhibitor (MEPI), protease inhibitor 14 (PI14), ZG-46p or serpin peptidase inhibitor clade I member 2 (serpin I2), is a protein belonging to the serpin family of serine cysteine protease inhibitors. Members of this superfamily play important roles in angiogenesis, inflammation, blood coagulation and apoptosis. Pancpin is expressed in pancreas and adipose tissues and localizes to the Golgi and zymogen granules. The down regulation of pancpin may be involved in the progression or development of pancreatic cancer. Overexpression of pancpin plays a significant role in inhibiting tumor formation. A mutation in the gene encoding pancpin can result in progressive apoptosis of pancreatic acinar cells. This can lead to pancreatic insufficiency; a disease characterized by an inability to absorb and digest nutrients from the diet.

## REFERENCES

- Chen, C.Y., Cronshagen, U. and Kern, H.F. 1997. A novel pancreas-specific serpin (ZG-46p) localizes to the soluble and membrane fraction of the Golgi complex and the zymogen granules of acinar cells. *Eur. J. Cell Biol.* 73: 205-214.
- Ozaki, K., Nagata, M., Suzuki, M., Fujiwara, T., Miyoshi, Y., Ishikawa, O., Ohigashi, H., Imaoka, S., Takahashi, E. and Nakamura, Y. 1998. Isolation and characterization of a novel human pancreas-specific gene, pancpin, that is down-regulated in pancreatic cancer cells. *Genes Chromosomes Cancer* 22: 179-185.
- Xiao, G., Liu, Y.E., Gentz, R., Sang, Q.A., Ni, J., Goldberg, I.D. and Shi, Y.E. 1999. Suppression of breast cancer growth and metastasis by a serpin myoepithelium-derived serine proteinase inhibitor expressed in the mammary myoepithelial cells. *Proc. Natl. Acad. Sci. USA* 96: 3700-3705.
- Chang, W.S., Chang, N.T., Lin, S.C., Wu, C.W. and Wu, F.Y. 2000. Tissue-specific cancer-related serpin gene cluster at human chromosome band 3q26. *Genes Chromosomes Cancer* 29: 240-255.
- Gudjonsson, T., Ronnov-Jessen, L., Villadsen, R., Rank, F., Bissell, M.J. and Petersen, O.W. 2002. Normal and tumor-derived myoepithelial cells differ in their ability to interact with luminal breast epithelial cells for polarity and basement membrane deposition. *J. Cell Sci.* 115: 39-50.
- Johnson-Pais, T.L., Nellierry, M.J., Ammerman, D.G., Pathmanathan, D., Bhatia, P., Buller, C.L., Leach, R.J. and Hansen, M.F. 2003. Determination of a minimal region of loss of heterozygosity on chromosome 18q21.33 in osteosarcoma. *Int. J. Cancer* 105: 285-288.
- Loftus, S.K., Cannons, J.L., Incao, A., Pak, E., Chen, A., Zerfas, P.M., Bryant, M.A., Biesecker, L.G., Schwartzberg, P.L. and Pavan, W.J. 2006. Acinar cell apoptosis in Serpini2-deficient mice models pancreatic insufficiency. *PLoS Genet.* 1: e38.

## CHROMOSOMAL LOCATION

Genetic locus: SERPINI2 (human) mapping to 3q26.1.

## PRODUCT

pancpin (h2): 293 Lysate represents a lysate of human pancpin transfected 293 cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

## APPLICATIONS

pancpin (h2): 293 Lysate is suitable as a Western Blotting positive control for human reactive pancpin antibodies. Recommended use: 10-20 µl per lane.

Control 293 Lysate: sc-110760 is available as a Western Blotting negative control lysate derived from non-transfected 293 cells.

## STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.