



**SZABO  
SCANDIC**

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](http://linkedin.com/company/szaboscandic)



# SAPK4 (h3): 293 Lysate: sc-158947

## BACKGROUND

Lipopolysaccharide has been shown to induce tyrosine phosphorylation of a unique protein, designated p38. p38 is a member of the MAP kinase family with features most closely resembling those of the *Saccharomyces cerevisiae* protein Hog1. p38 and Hog1 share a TGY phosphorylation sequence, whereas most other MAP kinase family proteins have a TEY sequence. A related protein, p38 $\beta$ , has been shown to phosphorylate ATF-2 at a 20-fold higher rate than p38, suggesting distinct substrate preferences. Stress activated protein kinase-4, or SAPK4, also designated p38 $\delta$ , is a related protein that is phosphorylated by MKK6 in response to cytokines and cellular stresses.

## REFERENCES

1. Han, J., Lee, J.D., Tobias, P.S. and Ulevitch, R.J. 1993. Endotoxin induces rapid protein tyrosine phosphorylation in 70Z/3 cells expressing CD14. *J. Biol. Chem.* 268: 25009-25014.
2. Brewster, J.L., de Valoir, T., Dwyer, N.D., Winter, E. and Gustin, M.C. 1993. An osmosensing signal transduction pathway in yeast. *Science* 259: 1760-1763.
3. Nishida, E. and Gotoh, Y. 1993. The MAP kinase cascade is essential for diverse signal transduction pathways. *Trends Biochem. Sci.* 18: 128-131.
4. Han, J., Lee, J.D., Bibbs, L. and Ulevitch, R.J. 1994. A MAP kinase targeted by endotoxin and hyperosmolarity in mammalian cells. *Science* 265: 808-811.
5. Jiang, Y., Chen, C., Li, Z., Guo, W., Gegner, J.A., Lin, S. and Han, J. 1996. Characterization of the structure and function of a new mitogen-activated protein kinase (p38 $\beta$ ). *J. Biol. Chem.* 271: 17920-17926.
6. Goedert, M., Cuenda, A., Craxton, M., Jakes, R. and Cohen, P. 1997. Activation of the novel stress-activated protein kinase SAPK4 by cytokines and cellular stresses is mediated by SKK3 (MKK6); comparison of its substrate specificity with that of other SAP kinases. *EMBO J.* 16: 3563-3571.
7. Kumar, S., McDonnell, P.C., Gum, R.J., Hand, A.T., Lee, J.C. and Young, P.R. 1997. Novel homologues of CSBP/p38 MAP kinase: activation, substrate specificity and sensitivity to inhibition by pyridinyl imidazoles. *Biochem. Biophys. Res. Commun.* 235: 533-538.
8. Wang, X.S., Diener, K., Manthey, C.L., Wang, S.W., Rosenweig, B., Bray, J., Delaney, J., Cole, C., Zukowski, M. and Yao, Z. 1997. Molecular cloning and characterization of a novel p38 mitogen activated protein kinase. *J. Biol. Chem.* 272: 23668-23674.

## CHROMOSOMAL LOCATION

Genetic locus: MAPK13 (human) mapping to 6p21.31.

## PRODUCT

SAPK4 (h3): 293 Lysate represents a lysate of human SAPK4 transfected 293 cells and is provided as 100  $\mu$ g protein in 200  $\mu$ l SDS-PAGE buffer.

## STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.


## APPLICATIONS

SAPK4 (h3): 293 Lysate is suitable as a Western Blotting positive control for human reactive SAPK4 antibodies. Recommended use: 10-20  $\mu$ l per lane.

Control 293 Lysate: sc-110760 is available as a Western Blotting negative control lysate derived from non-transfected 293 cells.

SAPK4 (20): sc-136063 is recommended as a positive control antibody for Western Blot analysis of enhanced human SAPK4 expression in SAPK4 transfected 293 cells (starting dilution 1:100, dilution range 1:100-1:1,000).

## DATA



SAPK4 (20): sc-136063. Western blot analysis of SAPK4 expression in non-transfected: sc-110760 (**A**) and human SAPK4 transfected: sc-158947 (**B**) 293 whole cell lysates.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.