



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Tenascin-X (h2): 293 Lysate: sc-159013

BACKGROUND

The tenascin family of extracellular matrix proteins includes tenascin (also designated cytotactin or tenascin-C) and tenascin-R (also designated restrictin or janusin) and tenascin-X. Tenascin proteins function as substrate-adhesion molecules (SAMs) and are involved in regulating numerous developmental processes, such as morphogenetic cell migration and organogenesis. The tenascin family proteins arise from various splicing events in the region of coding for FNIII repeats. Tenascin and tenascin-X are expressed in several tissues during embryogenesis, and in adult tissues undergoing active remodeling such as healing wounds and tumors. Tenascin-R (TN-R) is expressed on the surface of neurons and glial cells.

REFERENCES

1. Jung, M., Pesheva, P., Schachner, M. and Trotter, J. 1993. Astrocytes and neurons regulate the expression of the neural recognition molecule janusin by cultured oligodendrocytes. *Glia* 9: 163-175.
2. Schachner, M., Taylor, J., Bartsch, U. and Pesheva, P. 1994. The perplexing multifunctionality of janusin, a tenascin-related molecule. *Perspect. Dev. Neurobiol.* 2: 33-41.
3. Chiquet-Ehrismann, R. 1995. Tenascins, a growing family of extracellular matrix proteins. *Experientia* 51: 853-862.
4. Faissner, A. 1997. The tenascin gene family in axon growth and guidance. *Cell Tissue Res.* 290: 331-341.
5. Elefteriou, F., Exposito, J.Y., Garrone, R. and Lethias, C. 1997. Characterization of the bovine tenascin-X. *J. Biol. Chem.* 272: 22866-22874.
6. Srinivasan, J., Schachner, M. and Catterall, W.A. 1998. Interaction of voltage-gated sodium channels with the extracellular matrix molecules tenascin-C and tenascin-R. *Proc. Natl. Acad. Sci. USA* 95: 15753-15757.

CHROMOSOMAL LOCATION

Genetic locus: TNXB (human) mapping to 6p21.33.

PRODUCT

Tenascin-X (h2): 293 Lysate represents a lysate of human Tenascin-X transfected 293 cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

Tenascin-X (h2): 293 Lysate is suitable as a Western Blotting positive control for human reactive Tenascin-X antibodies. Recommended use: 10-20 µl per lane.

Control 293 Lysate: sc-110760 is available as a Western Blotting negative control lysate derived from non-transfected 293 cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.