



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



PSGL-1 (h3): 293 Lysate: sc-159212

BACKGROUND

PSGL-1, (P-Selectin glycoprotein ligand (also designated CD162) exists as a disulfide-linked homodimer. PSGL-1 is a type 1 membrane protein that localizes on the tips of microvilli of leukocytes. Its extracellular domain is rich in serines, threonines and prolines, and includes a series of 15 and 16 deca-meric repeats in HL-60 and U-937 cells, and human leukocytes, respectively. Although PSGL-1 appears to be the sole receptor for P-Selectin on human hematopoietic cells, it also interacts with E-Selectin through a unique binding site. In order to bind PSGL-1 to either E-Selectin or P-Selectin, PSGL-1 must be sialylated and fucosylated. PSGL-1 is a mucin-like molecule, much like leukosialin (CD43), CD164 and CD34. These proteins belong to an emerging family of cell adhesion receptors called sialomucins, which transduce negative signals in hematopoietic cells.

REFERENCES

1. Moore, K., Stults, N., Diaz, S., Smith, D., Cummings, R., Varki, A. and McEver, R. 1992. Identification of a specific glycoprotein ligand for P-Selectin (CD62) on myeloid cells. *J. Biol. Chem.* 118: 445-456.
2. Sako, D., Chang, X., Barone, K., Vachino, G., White, H., Shaw, G., Veldman, G., Bean, K., Ahern, T. and Furie, B. 1993. Expression cloning of a functional glycoprotein ligand for P-Selectin. *Cell* 75: 1179-1186.
3. Veldman, G., Bean, K., Cumming, D., Eddy, R. and Sait, S. 1995. Genomic organization and chromosomal localization of the gene encoding human P-Selectin glycoprotein ligand. *J. Biol. Chem.* 7: 16470-16475.
4. Patel, K., Moore, K., Nollert, M. and McEver, R. 1995. Neutrophils use both shared and distinct mechanisms to adhere to selectins under static and flow conditions. *J. Clin. Invest.* 96: 1887-1896.
5. Li, F., Erickson, H., James, J., Moore, K., Cummings, R. and McEver, R. 1996. Visualization of P-Selectin glycoprotein ligand-1 as a highly extended molecule and mapping of protein epitopes for monoclonal antibodies. *J. Biol. Chem.* 271: 6342-6348.
6. Levesque, J.P., Zannettino, A.C., Pudney, M., Niutta, S., Haylock, D., Snapp, K., Kansas, G., Berndt, M. and Simmons, P. 1999. PSGL-1-mediated adhesion of human hematopoietic progenitors to P-Selectin results in suppression of hematopoiesis. *Immunity* 11: 369-378.

CHROMOSOMAL LOCATION

Genetic locus: SELPLG (human) mapping to 12q24.11.

PRODUCT

PSGL-1 (h3): 293 Lysate represents a lysate of human PSGL-1 transfected 293 cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

PSGL-1 (h3): 293 Lysate is suitable as a Western Blotting positive control for human reactive PSGL-1 antibodies. Recommended use: 10-20 µl per lane.

Control 293 Lysate: sc-110760 is available as a Western Blotting negative control lysate derived from non-transfected 293 cells.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.