



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

NF-L (h): 293T Lysate: sc-159319

BACKGROUND

Neurofilament-L (NF-L), for neurofilament light polypeptide, a member of the intermediate filament family, is a major component of neuronal cytoskeletons. Neurofilaments are dynamic structures; they contain phosphorylation sites for a large number of protein kinases, including protein kinase A, protein kinase C, cyclin-dependent kinase 5, extracellular signal regulated kinase, glycogen synthase kinase-3 and stress-activated protein kinase γ . In addition to their role in the control of axon caliber, neurofilaments may affect other cytoskeletal elements, such as microtubules and Actin filaments. Changes in neurofilament phosphorylation or metabolism are frequently observed in neurodegenerative diseases, including amyotrophic lateral sclerosis (ALS), Parkinson's disease and Alzheimer's disease.

REFERENCES

- Angelides, K.J., et al. 1989. Assembly and exchange of intermediate filament proteins of neurons: neurofilaments are dynamic structures. *J. Cell Biol.* 108: 1495-1506.
- Sihag, R.K. and Nixon, R.A. 1989. *In vivo* phosphorylation of distinct domains of the 70-kilodalton neurofilament subunit involves different protein kinases. *J. Biol. Chem.* 264: 457-464.
- Hisanaga, S., et al. 1990. Effects of phosphorylation of the neurofilament L protein on filamentous structures. *Cell Regul.* 1: 237-248.
- Gonda, Y., et al. 1990. Involvement of protein kinase C in the regulation of assembly-disassembly of neurofilaments *in vitro*. *Biochem. Biophys. Res. Commun.* 167: 1316-1325.
- Nakamura, Y, et al. 1997. Abnormal distribution of neurofilament L in neurons with Alzheimer's disease. *Neurosci. Lett.* 225: 201-204.
- Hirokawa, N., et al. 1998. Gene targeting studies begin to reveal the function of neurofilament proteins. *J. Cell Biol.* 143: 1-4.
- Nakamura, Y., et al. 1999. Casein kinase II is responsible for phosphorylation of NF-L at Ser-473. *FEBS Lett.* 455: 83-86.
- Strong, M.J. 1999. Neurofilament metabolism in sporadic amyotrophic lateral sclerosis. *J. Neurol. Sci.* 169: 170-177.

CHROMOSOMAL LOCATION

Genetic locus: NEFL (human) mapping to 8p21.2.

PRODUCT

NF-L (h): 293T Lysate represents a lysate of human NF-L transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

APPLICATIONS

NF-L (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive NF-L antibodies. Recommended use: 10-20 μ l per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

NF-L (8A1): sc-20012 is recommended as a positive control antibody for Western Blot analysis of enhanced human NF-L expression in NF-L transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



NF-L (8A1): sc-20012. Western blot analysis of NF-L expression in non-transfected: sc-117752 (A) and human NF-L transfected: sc-159319 (B) 293T whole cell lysates.



NF-L (8A1): sc-20012. Western blot analysis of NF-L expression in non-transfected: sc-117752 (A) and human NF-L transfected: sc-159319 (B) 293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.