



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# Polycystin-L (h2): 293T Lysate: sc-159342

## BACKGROUND

Polycystin-L, also known as PKD2L1 (polycystic kidney disease 2-like 1), PCL, PKDL or PKD2L, is an 805 amino acid multi-pass membrane protein that belongs to the polycystin family of transmembrane proteins. Expressed in brain, heart, testis, spleen, liver and skeletal muscle, as well as in fetal kidney and liver, Polycystin-L functions as a calcium-regulated cation channel that is permeable to sodium, potassium and calcium and is involved in cell-cell and cell-matrix interactions. Polycystin-L shares 50% amino acid identity with a related family member, Polycystin-2, suggesting that Polycystin-L may be involved in the pathogenesis of polycystic kidney disease. Multiple alternatively spliced isoforms of Polycystin-L exist, all of which are encoded by a gene that maps to human chromosome 10q24.31.

## REFERENCES

1. Wu, G., Hayashi, T., Park, J.H., Dixit, M., Reynolds, D.M., Li, L., Maeda, Y., Cai, Y., Coca-Prados, M. and Somlo, S. 1998. Identification of PKD2L, a human PKD2-related gene: tissue-specific expression and mapping to chromosome 10q25. *Genomics* 54: 564-568.
2. Nomura, H., Turco, A.E., Pei, Y., Kalaydjieva, L., Schiavello, T., Weremowicz, S., Ji, W., Morton, C.C., Meisler, M., Reeders, S.T. and Zhou, J. 1998. Identification of PKDL, a novel polycystic kidney disease 2-like gene whose murine homologue is deleted in mice with kidney and retinal defects. *J. Biol. Chem.* 273: 25967-25973.
3. Guo, L., Chen, M., Basora, N. and Zhou, J. 2000. The human polycystic kidney disease 2-like (PKDL) gene: exon/intron structure and evidence for a novel splicing mechanism. *Mamm. Genome* 11: 46-50.
4. Stayner, C. and Zhou, J. 2001. Polycystin channels and kidney disease. *Trends Pharmacol. Sci.* 22: 543-546.
5. Li, Q., Liu, Y., Zhao, W. and Chen, X.Z. 2002. The calcium-binding EF-hand in Polycystin-L is not a domain for channel activation and ensuing inactivation. *FEBS Lett.* 516: 270-278.
6. Basora, N., Nomura, H., Berger, U.V., Stayner, C., Guo, L., Shen, X. and Zhou, J. 2002. Tissue and cellular localization of a novel polycystic kidney disease-like gene product, Polycystin-L. *J. Am. Soc. Nephrol.* 13: 293-301.
7. Li, Q., Liu, Y., Shen, P.Y., Dai, X.Q., Wang, S., Smillie, L.B., Sandford, R. and Chen, X.Z. 2003. Troponin I binds Polycystin-L and inhibits its calcium-induced channel activation. *Biochemistry* 42: 7618-7625.
8. Geng, L., Okuhara, D., Yu, Z., Tian, X., Cai, Y., Shibasaki, S. and Somlo, S. 2006. Polycystin-2 traffics to cilia independently of Polycystin-1 by using an N-terminal RVxP motif. *J. Cell Sci.* 119: 1383-1395.
9. Li, Q., Dai, X.Q., Shen, P.Y., Wu, Y., Long, W., Chen, C.X., Hussain, Z., Wang, S. and Chen, X.Z. 2007. Direct binding of  $\alpha$ -actinin enhances TRPP3 channel activity. *J. Neurochem.* 103: 2391-2400.

## CHROMOSOMAL LOCATION

Genetic locus: PKD2L1 (human) mapping to 10q24.31.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PRODUCT

Polycystin-L (h2): 293T Lysate represents a lysate of human Polycystin-L transfected 293T cells and is provided as 100  $\mu$ g protein in 200  $\mu$ l SDS-PAGE buffer.

## APPLICATIONS

Polycystin-L (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive Polycystin-L antibodies. Recommended use: 10-20  $\mu$ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

## STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.