

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



Inhibin β -A (h2): 293T Lysate: sc-159361



The Power to Ouestion

BACKGROUND

The $TGF\beta$ superfamily is composed of numerous growth and differentiation factors, including transforming growth factor β (TGF β) 1, 2 and 3; growth/ differentiation factor (GDF) 1 through 8; Mullerian inhibiting substance (MIS); bone morphogenic protein (BMP) 2 through 8; glial cell line-derived neurotrophic factor (GDNF); Inhibins (α , β -A, β -B and β -C), Lefty and Nodal. Members of the TGFβ superfamily are involved in embryonic development and adult tissue homeostasis. Inhibins and activins inhibit and activate, respectively, the secretion of follitropin by the pituitary gland. Inhibins and activins are involved in regulating a number of functions such as hypothalamic and pituitary hormone secretion, gonadal hormone secretion, germ cell development and maturation, erythroid differentiation, Insulin secretion, nerve cell survival, embryonic axial development or bone growth, depending on their subunit composition. Activins oppose the funtions of Inhibins. Inhibins are predominantly expressed in liver, uterus and ovary tissue. Inhibin A, a dimer of α and β -A, and Inhibin B, a dimer of α and β -B, have been shown to inhibit the secretion of follicle stimulating hormone. Inhibin β-C forms a homodimer and its expression is predominant in adult liver.

REFERENCES

- Stewart, A.G., et al. 1986. Human Inhibin genes. Genomic characterization and sequencing. FEBS Lett. 206: 329-334.
- 2. Mayo, K.E., et al. 1986. Inhibin A-subunit cDNAs from porcine ovary and human placenta. Proc. Natl. Acad. Sci. USA 83: 5849-5853.
- 3. Massague, J., et al. 1987. Multiple type- β transforming growth factors and their receptors. J. Cell. Physiol. Suppl. 5: 43-47.
- 4. Massague, J. 1990. The transforming growth factor- β family. Annu. Rev. Cell Biol. 6: 597-641.
- Albano, R.M., et al. 1993. Activins are expressed in preimplantation mouse embryos and in ES and EC cells and are regulated on their differentiation. Development 117: 711-723.
- 6. Schmitt, J., et al. 1996. Structure, chromosomal localization and expression analysis of the mouse Inhibin/Activin β C (Inhbc) gene. Genomics 32: 358-366.
- 7. McPherron, A.C., et al. 1997. Regulation of skeletal muscle mass in mice by a new TGF β superfamily member. Nature 387: 83-90.

CHROMOSOMAL LOCATION

Genetic locus: INHBA (human) mapping to 7p14.1.

PRODUCT

Inhibin β -A (h2): 293T Lysate represents a lysate of human Inhibin β -A transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

Inhibin β -A (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive Inhibin β -A antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3800 fax 831.457.3801 Europe +00800 4573 8000 49 6221 4503 0 www.scbt.com