



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# IFN- $\alpha$ / $\beta$ R $\beta$ (h2): 293T Lysate: sc-159501

## BACKGROUND

The type I interferons, IFN- $\alpha$  and IFN- $\beta$ , are a group of structurally and functionally related proteins that are induced by either viruses or double-stranded RNA and are defined by their ability to confer an antiviral state in cells. IFN- $\alpha$  and IFN- $\beta$  appear to compete with one another for binding to a common cell surface receptor, while immune IFN (IFN- $\gamma$ ) binds to a distinct receptor. This distinct receptor, IFN- $\alpha$ R, is only weakly responsive to type I interferons, in contrast to IFN- $\alpha$ / $\beta$ R, which binds to and responds effectively to IFN- $\beta$  and to several of the IFN- $\alpha$  subtypes. IFN- $\alpha$ / $\beta$ R is expressed as two alternatively spliced transcripts, designated IFN- $\alpha$ / $\beta$ R $\alpha$  (IFN- $\alpha$ / $\beta$ R1) and IFN- $\alpha$ / $\beta$ R $\beta$  (IFN- $\alpha$ / $\beta$ R2), both of which are involved in signal transduction and ligand binding.

## REFERENCES

1. Branca, A.A., et al. 1981. Evidence that type I and II interferons have different receptors. *Nature* 294: 768-770.
2. Orchansky, P., et al. 1984. Type I and type II interferon receptors. *J. Interferon Res.* 4: 275-282.
3. Novick, D., et al. 1987. The human interferon- $\gamma$  receptor, purification, characterization and preparation of antibodies. *J. Biol. Chem.* 262: 8483-8487.
4. Aguet, M., et al. 1988. Molecular cloning and expression of the human interferon- $\gamma$  receptor. *Cell* 55: 273-280.
5. Soh, J., et al. 1994. Identification and sequence of an accessory factor required for activation of the human interferon  $\gamma$  receptor. *Cell* 76: 793-802.
6. Hemmi, S., et al. 1994. A novel member of the interferon receptor family complements functionality of the murine interferon  $\gamma$  receptor in human cells. *Cell* 76: 803-810.
7. Novick, D., et al. 1994. The human interferon  $\alpha$ / $\beta$  receptor: characterization and molecular cloning. *Cell* 77: 391-400.
8. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 602376. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

## CHROMOSOMAL LOCATION

Genetic locus: IFNAR2 (human) mapping to 21q22.11.

## PRODUCT

IFN- $\alpha$ / $\beta$ R $\beta$  (h2): 293T Lysate represents a lysate of human IFN- $\alpha$ / $\beta$ R $\beta$  transfected 293T cells and is provided as 100  $\mu$ g protein in 200  $\mu$ l SDS-PAGE buffer.

## STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## APPLICATIONS

IFN- $\alpha$ / $\beta$ R $\beta$  (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive IFN- $\alpha$ / $\beta$ R $\beta$  antibodies. Recommended use: 10-20  $\mu$ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.