



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien


T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



L-Plastin (h3): 293T Lysate: sc-159508

BACKGROUND

Plastins (fimbrins) are members of a family of Actin-binding proteins that exhibit a tissue-specific expression pattern. Both L- and T-Plastin have been shown to be involved in cytoskeletal reorganization. L-Plastin, which is specifically expressed in hematopoietic cell lineages, has been proposed to be involved in the control of cell adhesion and motility. It is frequently expressed in cell lines derived from mammary solid tumors and is implicated in cancer invasion and metastasis. L-Plastin is also expressed in the majority of human cancer cell lines that are derived from various types of solid tumors. Additionally, L-Plastin is involved in regulating of leukocyte adhesion, and the phosphorylation of L-Plastin is implicated in modulating integrin regulation signaling pathways. T-Plastin is unique in that it is expressed in many types of tissues and notably absent in leukocytes.

REFERENCES

1. Lin, C.S., Lau, A. and Lue, T.F. 1998. Analysis and mapping of Plastin phosphorylation. *DNA Cell Biol.* 17: 1041-1046.
2. Jones, S.L., Wang, J., Turck, C.W. and Brown, E.J. 1998. A role for the Actin-bundling protein L-Plastin in the regulation of leukocyte integrin function. *Proc. Natl. Acad. Sci. USA* 95: 9331-9336.
3. Lin, C.S., Lau, A., Huynh, T. and Lue, T.F. 1999. Differential regulation of human T-Plastin gene in leukocytes and non-leukocytes: identification of the promoter, enhancer, and CpG island. *DNA Cell Biol.* 18: 27-37.
4. Lin, C.S., Lau, A., Yeh, C.C., Chang, C.H. and Lue, T.F. 2000. Upregulation of L-Plastin gene by testosterone in breast and prostate cancer cells: identification of three cooperative androgen receptor-binding sequences. *DNA Cell Biol.* 19: 1-7.
5. Lapillonne, A., Coue, O., Friederich, E., Nicolas, A., Del Maestro, L., Louvard, D., Robine, S. and Sastre-Garau, X. 2000. Expression patterns of L-Plastin isoform in normal and carcinomatous breast tissues. *Anticancer Res.* 20: 3177-3182.

CHROMOSOMAL LOCATION

Genetic locus: LCP1 (human) mapping to 13q14.13.

PRODUCT

L-Plastin (h3): 293T Lysate represents a lysate of human L-Plastin transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

L-Plastin (h3): 293T Lysate is suitable as a Western Blotting positive control for human reactive L-Plastin antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.