



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)

DNase I (h): 293T Lysate: sc-159797

BACKGROUND

Internucleosomal DNA fragmentation following the activation of endonucleases is the common end point of apoptosis. DNase I, a $\text{Ca}^{2+}/\text{Mg}^{2+}$ -dependent endonuclease ubiquitously expressed in mammalian tissues, has been implicated to mediate internucleosomal DNA degradation in human cells undergoing apoptosis. DNase I is highly polymorphic, and at least six alleles of DNase I are known. DNase II, the ubiquitously expressed acidic deoxyribonuclease, acts downstream of caspase activation and may also induce DNA digestion during apoptosis. DNase I cleaves DNA to 5'-phosphodinucleotide and 5'-phospho- oligonucleotide end-products, whereas DNase II cleaves DNA to 3'-phosphomononucleotide and 3'-phosphooligonucleotide end-products. The mechanism by which DNase II cuts DNA is similar to DNase I, which produces nicks rather than double-strand cuts. DNase II is usually present in cytoplasm of epithelial cells, but it appears concentrated in the nuclei of lens fibers. In contrast, DNase I is always concentrated in nuclei of epithelial and fiber cells. The gene encoding DNase II maps to human chromosome 19.

REFERENCES

1. Torriglia, A., Chaudun, E., Chany-Fournier, F., Jeanny, J.C., Courtois, Y. and Counis, M.F. 1995. Involvement of DNase II in nuclear degeneration during lens cell differentiation. *J. Biol. Chem.* 270: 28579-28585.
2. Yasuda, T., Takeshita, H., Iida, R., Nakajima, T., Hosomi, O., Nakashima, Y. and Kishi, K. 1998. Molecular cloning of the cDNA encoding human deoxyribonuclease II. *J. Biol. Chem.* 273: 2610-2616.
3. Krieser, R.J. and Eastman, A. 1998. The cloning and expression of human deoxyribonuclease II. A possible role in apoptosis. *J. Biol. Chem.* 273: 30909-30914.
4. Baker, K.P., Baron, W.F., Henzel, W.J. and Spencer, S.A. 1998. Molecular cloning and characterization of human and murine DNase II. *Gene* 215: 281-289.
5. Yasuda, T., Takeshita, H., Iida, R., Kogure, S. and Kishi, K. 1999. A new allele, DNASE1*6, of human deoxyribonuclease I polymorphism encodes an Arg to Cys substitution responsible for its instability. *Biochem. Biophys. Res. Commun.* 260: 280-283.
6. Oliveri, M., Daga, A., Cantoni, C., Lunardi, C., Millo, R. and Puccetti, A. 2001. DNase I mediates internucleosomal DNA degradation in human cells undergoing drug-induced apoptosis. *Eur. J. Immunol.* 31: 743-751.

CHROMOSOMAL LOCATION

Genetic Locus: DNASE1 (human) mapping to 16p13.3.

PRODUCT

DNase I (h): 293T Lysate represents a lysate of human DNase I transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

DNase I (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive DNase I antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.