



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PhLP (h2): 293T Lysate: sc-159877

BACKGROUND

Phosducin-like protein (PhLP, PDCL) is an ethanol-responsive modulator of heterotrimeric G proteins. The protein shares extensive amino acid sequence homology with phosducin (Phd), a phosphoprotein expressed in retina and pineal gland. Both PhLP and Phd regulate G protein signaling by binding to the $\beta\gamma$ subunits of G proteins. PhLP interacts with $G_{\beta\gamma}$ via a short C-terminal binding site. Additionally, PhLP acts as a substrate for GRK 2 phosphorylation at the same C-terminal binding site between residues 195 and 218. PhLPs may participate directly in the regulation of calcium-evoked exocytosis in adrenal medullary chromaffin cells. Glycosylated PhLP regulates opioid receptor function in mouse brain.

REFERENCES

1. Miles, M.F., Barhite, S., Sganga, M. and Elliott, M. 1993. Phosducin-like protein: an ethanol-responsive potential modulator of guanine nucleotide-binding protein function. *Proc. Natl. Acad. Sci. USA* 90: 10831-10835.
2. Schroder, S., Bluml, K., Dees, C. and Lohse, M.J. 1997. Identification of a C-terminal binding site for G protein $\beta\gamma$ subunits in phosducin-like protein. *FEBS Lett.* 401: 243-246.
3. Thibault, C., Feng Wang, J., Charnas, R., Mirel, D., Barhite, S. and Miles, M.F. 1999. Cloning and characterization of the rat and human phosducin-like protein genes: structure, expression and chromosomal localization. *Biochim. Biophys. Acta* 1444: 346-354.
4. Ruiz-Gomez, A., Humrich, J., Murga, C., Quitterer, U., Lohse, M.J. and Mayor, F., Jr. 2000. Phosphorylation of phosducin and phosducin-like protein by G protein-coupled receptor kinase 2. *J. Biol. Chem.* 275: 29724-29730.
5. Gensse, M., Vitale, N., Chasserot-Golaz, S. and Bader, M.F. 2000. Regulation of exocytosis in chromaffin cells by phosducin-like protein, a protein interacting with G protein $\beta\gamma$ subunits. *FEBS Lett.* 480: 184-188.
6. Garzon, J., Rodriguez-Diaz, M., Lopez-Fando, A., Garcia-Espana, A. and Sanchez-Blazquez, P. 2002. Glycosylated phosducin-like protein long regulates opioid receptor function in mouse brain. *Neuropharmacology* 42: 813-828.

CHROMOSOMAL LOCATION

Genetic locus: PDCL (human) mapping to 9q33.2.

PRODUCT

PhLP (h): 293T Lysate represents a lysate of human PhLP transfected 293T cells and is provided as 100 μg protein in 200 μl SDS-PAGE buffer.

STORAGE

Store at -20°C . Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

APPLICATIONS

PhLP (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive PhLP antibodies. Recommended use: 10-20 μl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.