



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



NIRF (h): 293 Lysate: sc-170276

BACKGROUND

NIRF (Np95/ICBP90-like RING finger protein), also known as E3 ubiquitin-protein ligase UHRF2, nuclear zinc finger protein Np97 or RING finger protein 107, is a nuclear protein involved in cell cycle regulation. NIRF contains a PHD finger, two RING fingers, a ubiquitin-like domain and a YDG/SRA domain. It shares high structural homology with UHRF1 (also called ICBP90 in humans and Np95 in mice), however, in contrast to UHRF1, NIRF acts as a negative regulator of cell proliferation. It associates with the Cdk2-cyclin complex in its dephosphorylated form and induces G₁ arrest. NIRF plays an important role in the regulation of the G₁/S transition by blocking cell entry into the S phase. While associated with Cdk2, NIRF becomes phosphorylated. NIRF can also act as a ubiquitin ligase and it ubiquitinates PCNP. In addition, NIRF can recruit and bind HDAC1 via its SRA domain. The overexpression of NIRF results in an increase of G₁ phase cells.

REFERENCES

1. Mori, T., Li, Y., Hata, H., Ono, K. and Kochi, H. 2002. NIRF, a novel RING finger protein, is involved in cell-cycle regulation. *Biochem. Biophys. Res. Commun.* 296: 530-536.
2. Li, Y., Mori, T., Hata, H., Homma, Y. and Kochi, H. 2004. NIRF induces G₁ arrest and associates with Cdk2. *Biochem. Biophys. Res. Commun.* 319: 464-468.
3. Mori, T., Li, Y., Hata, H. and Kochi, H. 2004. NIRF is a ubiquitin ligase that is capable of ubiquitinating PCNP, a PEST-containing nuclear protein. *FEBS Lett.* 557: 209-214.
4. Unoki, M., Nishidate, T. and Nakamura, Y. 2004. ICBP90, an E2F-1 target, recruits HDAC1 and binds to methyl-CpG through its SRA domain. *Oncogene* 23: 7601-7610.
5. Abbady, A.Q., Bronner, C., Bathami, K., Muller, C.D., Jeanblanc, M., Mathieu, E., Klein, J.P., Candolfi, E. and Mousli, M. 2005. TCR pathway involves ICBP90 gene down-regulation via E2F binding sites. *Biochem. Pharmacol.* 70: 570-579.
6. Muto, M., Fujimori, A., Neno, M., Daino, K., Matsuda, Y., Kuroiwa, A., Kubo, E., Kanari, Y., Utsuno, M., Tsuji, H., Ukai, H., Mita, K., Takahagi, M. and Tatsumi, K. 2006. Isolation and Characterization of a novel human radiosusceptibility gene, NP95. *Radiat. Res.* 166: 723-733.
7. Woo, H.R., Pontes, O., Pikaard, C.S. and Richards, E.J. 2007. VIM1, a methylcytosine-binding protein required for centromeric heterochromatinization. *Genes Dev.* 21: 267-277.

CHROMOSOMAL LOCATION

Genetic locus: UHRF2 (human) mapping to 9p24.1.

PRODUCT

NIRF (h): 293 Lysate represents a lysate of human NIRF transfected 293 cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

NIRF (h): 293 Lysate is suitable as a Western Blotting positive control for human reactive NIRF antibodies. Recommended use: 10-20 µl per lane.

Control 293 Lysate: sc-110760 is available as a Western Blotting negative control lysate derived from non-transfected 293 cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.