



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

ADSL (h3): 293T Lysate: sc-170309

BACKGROUND

ADSL (adenylosuccinate lyase), also known as AMPS, ASL or ASASE, is a 484 amino acid protein that is involved in both purine biosynthesis and in the formation of adenosine monophosphate (AMP) from inosine monophosphate. Expressed ubiquitously, ADSL catalyzes two key reactions in AMP biosynthesis, namely the removal of a fumarate from succinylaminoimidazole carboxamide (SAICA) ribotide to give aminoimidazole carboxamide ribotide (AICA) and the subsequent removal of fumarate from adenylosuccinate to yield AMP. Defects in the gene encoding ADSL are the cause of adenylosuccinase deficiency (ADSL deficiency), an autosomal recessive disorder characterized by epilepsy, growth retardation and muscular wasting. Multiple isoforms of ADSL exist due to alternative splicing events.

REFERENCES

1. Kmoch, S., Hartmannová, H., Stiburková, B., Krijt, J., Zikánová, M. and Sebesta, I. 2000. Human adenylosuccinate lyase (ADSL), cloning and characterization of full-length cDNA and its isoform, gene structure and molecular basis for ADSL deficiency in six patients. *Hum. Mol. Genet.* 9: 1501-1513.
2. Race, V., Marie, S., Vincent, M.F. and Van den Berghe, G. 2000. Clinical, biochemical and molecular genetic correlations in adenylosuccinate lyase deficiency. *Hum. Mol. Genet.* 9: 2159-2165.
3. Tabucchi, A., Carlucci, F., Rosi, F., Guerranti, R. and Marinello, E. 2001. Determination, activity and biological role of adenylosuccinate lyase in blood cells. *Biomed. Pharmacother.* 55: 277-283.
4. Marie, S., Race, V., Nassogne, M.C., Vincent, M.F. and Van den Berghe, G. 2002. Mutation of a nuclear respiratory factor 2 binding site in the 5' untranslated region of the ADSL gene in three patients with adenylosuccinate lyase deficiency. *Am. J. Hum. Genet.* 71: 14-21.
5. Edery, P., Chabrier, S., Ceballos-Picot, I., Marie, S., Vincent, M.F. and Tardieu, M. 2003. Intrafamilial variability in the phenotypic expression of adenylosuccinate lyase deficiency: a report on three patients. *Am. J. Med. Genet. A* 120A: 185-190.
6. Sivendran, S., Patterson, D., Spiegel, E., McGown, I., Cowley, D. and Colman, R.F. 2004. Two novel mutant human adenylosuccinate lyases (ASLs) associated with autism and characterization of the equivalent mutant *Bacillus subtilis* ASL. *J. Biol. Chem.* 279: 53789-53797.
7. Lee, P. and Colman, R.F. 2007. Expression, purification, and characterization of stable, recombinant human adenylosuccinate lyase. *Protein Expr. Purif.* 51: 227-234.
8. Online Mendelian Inheritance in Man, OMIM™. 2009. Johns Hopkins University, Baltimore, MD. MIM Number: 608222. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: ADSL (human) mapping to 22q13.1.

PRODUCT

ADSL (h3): 293T Lysate represents a lysate of human ADSL transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

ADSL (h3): 293T Lysate is suitable as a Western Blotting positive control for human reactive ADSL antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

ADSL (C-11): sc-365623 is recommended as a positive control antibody for Western Blot analysis of enhanced human ADSL expression in ADSL transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



ADSL (C-11): sc-365623. Western blot analysis of ADSL expression in non-transfected: sc-117752 (A) and human ADSL transfected: sc-170309 (B) 293T whole cell lysates.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.