



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



Tuftelin (h3): 293T Lysate: sc-170311

BACKGROUND

Dental enamel is a highly mineralized tissue in which most of the volume is occupied by large, highly organized hydroxyapatite crystals. This structure is thought to be controlled through the interaction of many organic matrix molecules, including Amelogenin, Ameloblastin, Enamelin and Tuftelin. All of these secreted proteins are involved in the mineralization and enamel matrix formation in developing tooth enamel. Tuftelin is also expressed in normal and cancerous non-mineralizing soft tissues, which suggests it has a universal function and/or a multifunctional role. The Tuftelin protein contains one N-glycosylation site, seven O-glycosylation sites and seven phosphorylation sites. It also contains a coiled-coil domain that is involved in self-assembly and the interaction of Tuftelin with the Tuftelin interacting protein TIP39.

REFERENCES

1. Paine, M.L., Deutsch, D. and Snead, M.L. 1997. Carboxyl-region of Tuftelin mediates self-assembly. *Connect. Tissue Res.* 35: 157-161.
2. Deutsch, D., Dafni, L., Palmon, A., Hekmati, M., Young, M.F. and Fisher, L.W. 1997. Tuftelin: enamel mineralization and amelogenesis imperfecta. *Ciba Found. Symp.* 205: 135-147.
3. MacDougall, M., Simmons, D., Dodds, A., Knight, C., Luan, X., Zeichner-David, M., Zhang, C., Ryu, O.H., Qian, Q., Simmer, J.P. and Hu, C.C. 1998. Cloning, characterization, and tissue expression pattern of mouse Tuftelin cDNA. *J. Dent. Res.* 77: 1970-1978.
4. Deutsch, D., Palmon, A., Dafni, L., Mao, Z., Leytin, V., Young, M. and Fisher, L.W. 1998. Tuftelin—aspects of protein and gene structure. *Eur. J. Oral Sci.* 106: 315-323.
5. Mao, Z., Shay, B., Hekmati, M., Fermon, E., Taylor, A., Dafni, L., Heikinheimo, K., Lustmann, J., Fisher, L.W., Young, M.F. and Deutsch, D. 2001. The human Tuftelin gene: cloning and characterization. *Gene* 279: 181-196.
6. Deutsch, D., Leiser, Y., Shay, B., Fermon, E., Taylor, A., Rosenfeld, E., Dafni, L., Charuvi, K., Cohen, Y., Haze, A., Fuks, A. and Mao, Z. 2002. The human Tuftelin gene and the expression of Tuftelin in mineralizing and nonmineralizing tissues. *Connect. Tissue Res.* 43: 425-434.
7. Luo, W., Wen, X., Wang, H.J., MacDougall, M., Snead, M.L. and Paine, M.L. 2004. *In vivo* overexpression of Tuftelin in the enamel organic matrix. *Cells Tissues Organs* 177: 212-220.

CHROMOSOMAL LOCATION

Genetic locus: TUFT1 (human) mapping to 1q21.3.

PRODUCT

Tuftelin (h3): 293T Lysate represents a lysate of human Tuftelin transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

Tuftelin (h3): 293T Lysate is suitable as a Western Blotting positive control for human reactive Tuftelin antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.