



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# Moesin (h2): 293T Lysate: sc-170446

## BACKGROUND

Ezrin, Moesin and Radixin belong to a family of highly homologous Actin-associated proteins that are localized just beneath the plasma membrane. These proteins are believed to be involved in the mediation of interactions between cytoskeletal and membrane proteins. Ezrin serves as a major cytoplasmic substrate of various protein-tyrosine kinases, including the epidermal growth factor receptor. Ezrin has also been identified as a cAMP-dependent protein kinase (A-kinase) anchoring protein and designated AKAP78. Moesin and Radixin share more than 70% homology with Ezrin and are coexpressed within various cell types. Despite the high degree of homology, the three proteins exhibit a distinct receptor-specific pattern of phosphorylation.

## REFERENCES

- Gould, K.L., et al. 1989. cDNA cloning and sequencing of the protein-tyrosine kinase substrate, Ezrin, reveals homology to band 4.1. *EMBO J.* 8: 4133-4142.
- Lankes, W.T. and Furthmayr, H. 1991. Moesin: a member of the protein 4.1-Talin-Ezrin family of protein. *Proc. Natl. Acad. Sci. USA* 88: 8297-8301.
- Sato, N., et al. 1992. A gene family consisting of Ezrin, Radixin and Moesin. Its specific localization at Actin filament/plasma membrane association sites. *J. Cell Sci.* 103: 131-143.
- Fazioli, F., et al. 1993. The Ezrin-like family of tyrosine kinase substrates: receptor-specific pattern of tyrosine phosphorylation and relationship to malignant transformation. *Oncogene* 8: 1335-1345.
- Algrain, M., et al. 1993. Ezrin contains cytoskeleton and membrane binding domains accounting for its proposed role as a membrane-cytoskeletal linker. *J. Cell Biol.* 120: 129-139.
- Tsukita, S., et al. 1994. ERM family members as molecular linkers between the cell surface glycoprotein CD44 and Actin-based cytoskeletons. *J. Cell Biol.* 126: 391-401.
- Andreoli, C., et al. 1994. Ezrin has properties to self-associate at the plasma membrane. *J. Cell Sci.* 107: 2509-2521.

## CHROMOSOMAL LOCATION

Genetic locus: MSN (human) mapping to Xq12.

## PRODUCT

Moesin (h2): 293T Lysate represents a lysate of human Moesin transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

## APPLICATIONS

Moesin (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive Moesin antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

Moesin (38/87): sc-58806 is recommended as a positive control antibody for Western Blot analysis of enhanced human Moesin expression in Moesin transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:  
1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

## DATA



Moesin (38/87): sc-58806. Western blot analysis of Moesin expression in non-transfected: sc-117752 (A) and human Moesin transfected: sc-170446 (B) 293T whole cell lysates.

## STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.