



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



GI Syn (h2): 293T Lysate: sc-170514

BACKGROUND

Glutamine synthetase (GI Syn) forms a homo-octamer that serves as a catalyst for the amination of glutamic acid to form glutamine. This enzyme is a marker for astrocytes, which serve as the primary site of conversion of glutamic acid to glutamine in the brain. Induction of GI Syn is seen upon astrocyte cell contact with neurons. Elevated expression of GI Syn in glial cells has been shown to protect neurons from degeneration due to excess glutamate. GI Syn is also present in the liver and is involved in nitrogen homeostasis. Overexpression of GI Syn has been shown in primary liver cancers, indicating a potential role for GI Syn in hepatocyte transformation.

REFERENCES

1. Gibbs, C.S., Campbell, K.E. and Wilson, R.H. 1987. Sequence of a human glutamine synthetase cDNA. *Nucleic Acids Res.* 15: 6293.
2. Linser, P.J. and Perkins, M. 1987. Gliogenesis in the embryonic avian optic tectum: neuronal-glial interactions influence astroglial phenotype maturation. *Brain Res.* 428: 277-290.
3. Vardimon, L., Fox, L.L., Degenstein, L. and Moscona, A.A. 1988. Cell contacts are required for induction by cortisol of glutamine synthetase gene transcription in the retina. *Proc. Natl. Acad. Sci. USA* 85: 5981-5985.
4. Mill, J.F., Mearow, K.M., Purohit, H.J., Haleem-Smith, H., King, R. and Freese, E. 1991. Cloning and functional characterization of the rat glutamine synthetase gene. *Brain Res. Mol. Brain Res.* 9: 197-207.
5. Van den Hoff, M.J., Geerts, W.J., Das, A.T., Moorman, A.F. and Lamers, W.H. 1991. cDNA sequence of the long mRNA for human glutamine synthase. *Biochim. Biophys. Acta* 1090: 249-251.
6. Christa, L., Simon, M.T., Flinois, J.P., Gebhardt, R., Brechot, C. and Lasserre, C. 1994. Overexpression of glutamine synthetase in human primary liver cancer. *Gastroenterology* 106: 1312-1320.
7. Gorovits, R., Avidan, N., Avisar, N., Shaked, I. and Vardimon, L. 1997. Glutamine synthetase protects against neuronal degeneration in injured retinal tissue. *Proc. Natl. Acad. Sci. USA* 94: 7024-7029.

CHROMOSOMAL LOCATION

Genetic locus: GLUL (mouse) mapping to 1q25.3.

PRODUCT

GI Syn (h2): 293T Lysate represents a lysate of human GI Syn transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

GI Syn (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive GI Syn antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.