



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



TyrRS (h2): 293T Lysate: sc-170741

BACKGROUND

Tryptophanyl-tRNA synthetases are essential enzymes that catalyze the aminoacylation of tRNA^{Trp} with tryptophan, an essential function of the cell's protein synthesis machinery. Two forms of tryptophanyl-tRNA synthetase exist; a cytoplasmic form, named TrpRS (also known as WARS), and a mitochondrial form, named WARS2. In normal cells, human TrpRS exists as a full length form and as a truncated form designated mini TrpRS, which is produced by alternative splicing. Expression of mini TrpRS is highly stimulated in human cells by the addition of IFN- γ . Although both human full length TrpRS and mini TrpRS are enzymatically active in aminoacylation, they differ in angiostatic activity. The tyrosyl-tRNA synthetase (TyrRS), a protein related to TrpRS, exists as a homodimeric enzyme that catalyzes the aminoacylation of tRNA^{Tyr} by L-tyrosine.

REFERENCES

- Wakasugi, K., Slike, B.M., Hood, J., Otani, A., Ewalt, K.L., Friedlander, M., Cheresh, D.A. and Schimmel, P. 2002. A human aminoacyl-tRNA synthetase as a regulator of angiogenesis. Proc. Natl. Acad. Sci. USA 99: 173-177.
- Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 191050. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Jia, J., Li, B., Jin, Y. and Wang, D. 2003. Expression, purification, and characterization of human tyrosyl-tRNA synthetase. Protein Expr. Purif. 27: 104-108.
- Yang, X.L., Schimmel, P. and Ewalt, K.L. 2004. Relationship of two human tRNA synthetases used in cell signaling. Trends Biochem. Sci. 29: 250-256.
- Kise, Y., Lee, S.W., Park, S.G., Fukai, S., Sengoku, T., Ishii, R., Yokoyama, S., Kim, S. and Nureki, O. 2004. A short peptide insertion crucial for angiostatic activity of human tryptophanyl-tRNA synthetase. Nat. Struct. Mol. Biol. 11: 149-156
- Bonnefond, L., Frugier, M., Giege, R. and Rudinger-Thirion, J. 2005. Human mitochondrial TyrRS disobeys the tyrosine identity rules. RNA 11: 558-562.
- Bonnefond, L., Fender, A., Rudinger-Thirion, J., Giege, R., Florentz, C. and Sissler, M. 2005. Toward the full set of human mitochondrial aminoacyl-tRNA synthetases: characterization of AspRS and TyrRS. Biochemistry 44: 4805-4816.
- LocusLink Report (LocusID: 8565). <http://www.ncbi.nlm.nih.gov/LocusLink/>

CHROMOSOMAL LOCATION

Genetic locus: YARS (human) mapping to 1p35.1.

PRODUCT

TyrRS (h2): 293T Lysate represents a lysate of human TyrRS transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

TyrRS (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive TyrRS antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.