



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



mSHMT (h3): 293T Lysate: sc-170754



The Power to Question

BACKGROUND

Mammalian serine hydroxymethyltransferase (SHMT) is a tetrameric, pyridoxal phosphate (PLP)-dependent enzyme that catalyzes the reversible interconversion of serine and tetrahydrofolate to glycine and methylenetetrahydrofolate in the cytoplasm (cSHMT, SHMT1) and mitochondria (mSHMT, SHMT2). cSHMT preferentially supplies one-carbon units for thymidylate biosynthesis, depletes methylenetetrahydrofolate pools for S-adenosylmethionine (SAM) synthesis by synthesizing serine, sequesters 5-methyltetrahydrofolate, and inhibits SAM synthesis. Sheep liver cytosolic recombinant SHMT (scSHMT), Lys-71, Arg-80 and Asp-89 residues influence intra-subunit ionic interactions essential for catalytic activity; Tyr 72, Asp-227 and His-356 residues in the active site interact with PLP and maintain the tetrameric structure. The cDNA for the mitochondrial enzyme encodes a mature protein of 474 residues.

REFERENCES

1. Online Mendelian Inheritance in Man, OMIM™. 1998. Johns Hopkins University, Baltimore, MD. MIM Number: 138450. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
2. Liu, X., Szebenyi, D.M., Anguera, M.C., Thiel, D.J. and Stover, P.J. 2001. Lack of catalytic activity of a murine mRNA cytoplasmic serine hydroxymethyltransferase splice variant: evidence against alternative splicing as a regulatory mechanism. *Biochemistry* 40: 4932-4939.
3. Trivedi, V., Gupta, A., Jala, V.R., Saravanan, P., Rao, G.S., Rao, N.A., Savithri, H.S. and Subramanya, H.S. 2002. Crystal structure of binary and ternary complexes of serine hydroxymethyltransferase from *Bacillus stearothermophilus*: insights into the catalytic mechanism. *J. Biol. Chem.* 277: 17161-17169.
4. Herbig, K., Chiang, E.P., Lee, L.R., Hills, J., Shane, B. and Stover, P.J. 2002. Cytoplasmic serine hydroxymethyltransferase mediates competition between folate-dependent deoxyribonucleotide and S-adenosylmethionine biosyntheses. *J. Biol. Chem.* 277: 38381-38389.
5. Jala, V.R., Appaji Rao, N. and Savithri, H.S. 2004. Identification of amino acid residues, essential for maintaining the tetrameric structure of sheep liver cytosolic serine hydroxy-methyltransferase, by targeted mutagenesis. *Biochem. J.* 369: 469-476.
6. LocusLink Report (LocusID: 6472). <http://www.ncbi.nlm.nih.gov/LocusLink/>

CHROMOSOMAL LOCATION

Genetic locus: SHMT2 (human) mapping to 12q13.3.

PRODUCT

mSHMT (h3): 293T Lysate represents a lysate of human mSHMT transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

mSHMT (h3): 293T Lysate is suitable as a Western Blotting positive control for human reactive mSHMT antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.