



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



Rad1 (h2): 293T Lysate: sc-171349

BACKGROUND

DNA damage or incomplete replication of DNA results in inhibition of cell cycle progression at the G₁/S or G₂/M checkpoints by conserved regulatory mechanisms. Rad17 is involved in regulation of cell cycle arrest at the G₁ checkpoint, whereas Chk1, Rad1, Rad9 and Hus1 are involved in regulation of cell cycle arrest at the G₂ checkpoint. Overexpression of Rad17 results in p53 activation and an accumulation of cells in G₁ phase. Chk1 functions as an essential component in the G₂ DNA damage checkpoint by phosphorylating Cdc25C in response to DNA damage, thus inhibiting mitosis. Hus1 and Rad9 exhibit conserved function in fission yeast and higher eukaryotes. Hus1 has been shown to be phosphorylated in response to DNA damage, a process which requires Rad checkpoint genes. Rad9 is thought to be a candidate tumor suppressor gene because it is localized to a region of human chromosome 11, which is a region containing a number of tumor suppressor loci.

REFERENCES

1. Carr, A.M., Moudjou, M., Bentley, N.J. and Hagan, I.M. 1995. The Chk1 pathway is required to prevent mitosis following cell-cycle arrest at "start". *Curr. Biol.* 5: 1179-1190.
2. Lieberman, H.B., Hopkins, K.M., Nass, M., Demetrick, D. and Davey, S. 1996. A human homolog of the *Schizosaccharomyces pombe* Rad9⁺ checkpoint control gene. *Proc. Natl. Acad. Sci. USA* 93: 13890-13895.
3. Sanchez, Y., Wong, C., Thoma, R.S., Richman, R., Wu, Z., Piwnica-Worms, H. and Elledge, S.J. 1997. Conservation of the Chk1 checkpoint pathway in mammals: linkage of DNA damage to Cdk regulation through Cdc25. *Science* 277: 1497-1501.
4. Peng, C.Y., Graves, P.R., Thoma, R.S., Wu, Z., Shaw, A.S. and Piwnica-Worms, H. 1997. Mitotic and G₂ checkpoint control: regulation of 14-3-3 protein binding by phosphorylation of Cdc25C on Serine 216. *Science* 277: 1501-1505.
5. O'Connell, M.J., Raleigh, J.M., Verkade, H.M. and Nurse, P. 1997. Chk1 is a Wee 1 kinase in the G₂ DNA damage checkpoint inhibiting Cdc2 by Y15 phosphorylation. *EMBO J.* 16: 545-554.
6. Kostrub, C.F., Knudsen, K., Subramani, S. and Enoch, T. 1998. Hus1p, a conserved fission yeast checkpoint protein, interacts with Rad1p and is phosphorylated in response to DNA damage. *EMBO J.* 17: 2055-2066.
7. Bao, S., Chang, M.S., Auclair, D., Sun, Y., Wang, Y., Wong, W.K., Zhang, J., Liu, Y., Qian, X., Sutherland, R., Magi-Galluzzi, C., Weisberg, E., Cheng, E.Y., Hao, L., Sasaki, H., Campbell, M.S., Kraeft, S.K., Loda, M., Lo, K.M. and Chen, L.B. 1999. HRAD17, a human homologue of the *Schizosaccharomyces pombe* checkpoint gene rad17, is overexpressed in colon carcinoma. *Cancer Res.* 59: 2023-2028.
8. Nicholson, A., Fabbri, R.M., Reeves, J.W. and Crouse, G.F. 2006. The effects of mismatch repair and RAD1 genes on interchromosomal crossover recombination in *Saccharomyces cerevisiae*. *Genetics* 173: 647-659.
9. Francia, S., Weiss, R.S. and d'Adda di Fagagna, F. 2007. Need telomere maintenance? Call 911. *Cell Div.* 2: 3.

CHROMOSOMAL LOCATION

Genetic locus: RAD1 (human) mapping to 5p13.2.

PRODUCT

Rad1 (h2): 293T Lysate represents a lysate of human Rad1 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

Rad1 (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive Rad1 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.