



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)

# ASCT1 (h3): 293T Lysate: sc-171350

## BACKGROUND

Neutral amino acid transporter proteins, also designated alanine/serine/cysteine/threonine transporters (ASCT), belong to the sodium:dicarboxylate (SDF) symporter family or proteins. The members of this family of proteins are multi-pass membrane-bound proteins that act as transporters for threonine, alanine, serine and cysteine. ASCT1 and ASCT2 have been shown to exhibit sodium dependence. ASCT1 is expressed in most tissues, but highest expression has been detected in muscle, brain and pancreas. The highest levels of ASCT2 expression are found in placenta, kidney, pancreas, muscle and intestine.

## REFERENCES

1. Arriza, J.L., et al. 1993. Cloning and expression of a human neutral amino acid transporter with structural similarity to the glutamate transporter gene family. *J. Biol. Chem.* 268: 15329-15332.
2. Hofmann, K., et al. 1995. Human neutral amino acid transporter ASCT1: structure of the gene (SLC1A4) and localization to chromosome 2p13-p15. *Genomics* 24: 20-26.
3. Kekuda, R., et al. 1996. Cloning of the sodium-dependent, broad-scope, neutral amino acid transporter Bo from a human placental choriocarcinoma cell line. *J. Biol. Chem.* 271: 18657-18661.
4. Rasko, J.E., et al. 1999. The RD114/simian type D retrovirus receptor is a neutral amino acid transporter. *Proc. Nat. Acad. Sci. USA* 96: 2129-2134.
5. Tailor, C.S., et al. 1999. A sodium-dependent neutral-amino-acid transporter mediates infections of feline and baboon endogenous retroviruses and simian type D retroviruses. *J. Virol.* 73: 4470-4474.
6. Tailor, C.S., et al. 2001. Truncated forms of the dual function human ASCT2 neutral amino acid transporter/retroviral receptor are translationally initiated at multiple alternative CUG and GUG codons. *J. Biol. Chem.* 276: 27221-27230.
7. Yamamoto, T., et al. 2004. Functional identification of ASCT1 neutral amino acid transporter as the predominant system for the uptake of L-serine in rat neurons in primary culture. *Neurosci. Res.* 49: 101-111.
8. Kanai, Y. and Hediger, M.A. 2004. The glutamate/neutral amino acid transporter family SLC1: molecular, physiological and pharmacological aspects. *Pflugers Arch.* 447: 469-479.
9. Hashimoto, Y., et al. 2005. Distribution of neutral amino acid transporter ASCT1 in the non-neuronal tissues of mice. *Jpn. J. Vet. Res.* 52: 113-124.

## CHROMOSOMAL LOCATION

Genetic locus: SLC1A4 (human) mapping to 2p14.

## PRODUCT

ASCT1 (h3): 293T Lysate represents a lysate of human ASCT1 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## APPLICATIONS

ASCT1 (h3): 293T Lysate is suitable as a Western Blotting positive control for human reactive ASCT1 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

## STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.