



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



PPIL1 (h): 293T Lysate: sc-171444

BACKGROUND

PPIL1 (peptidylprolyl isomerase (cyclophilin)-like 1), also known as CYPL1, PPIase, CGI-124 or hCypX, is a member of the cyclophilin-type PPIase family of proteins. PPIL1 contains one PPIase cyclophilin-type domain and is ubiquitously expressed with predominant expression in skeletal muscle and heart. PPIL1 is a component of the 35 S U5 snRNP (small nuclear ribonucleoprotein) and is also recruited to the 45 S activated spliceosome by Skip (SNW1), a transcriptional coactivator. PPIL1 stably associates with Skip and may play a role in spliceosome activation, possibly functioning as a foldase or a molecular chaperone. In addition, PPIL1 interacts with Op18, a protein involved in microtubule stabilization, and may participate in cell proliferation. PPIL1 expression levels are elevated in cancer cells, further supporting a role for PPIL1 in proliferation and tumorigenesis.

REFERENCES

1. Ozaki, K., Fujiwara, T., Kawai, A., Shimizu, F., Takami, S., Okuno, S., Takeda, S., Shimada, Y., Nagata, M., Watanabe, T., Takaichi, A., Takahashi, E., Nakamura, Y. and Shin, S. 1996. Cloning, expression and chromosomal mapping of a novel cyclophilin-related gene (PPIL1) from human fetal brain. *Cytogenet. Cell Genet.* 72: 242-245.
2. Mann, S.S., Pettenati, M.J., von Kap-herr, C. and Hart, T.C. 1998. Reassignment of peptidyl prolyl isomerase-like 1 gene (PPIL1) to human chromosome region 6p21.1 by radiation hybrid mapping and fluorescence *in situ* hybridization. *Cytogenet. Cell Genet.* 83: 228-229.
3. Skruzny, M., Ambrozkova, M., Fukova, I., Martinkova, K., Blahuskova, A., Hamplova, L., Puta, F. and Folk, P. 2001. Cyclophilins of a novel subfamily interact with SNW/SKIP coregulator in *Dictyostelium discoideum* and *Schizosaccharomyces pombe*. *Biochim. Biophys. Acta* 1521: 146-151.
4. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 601301. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
5. Folk, P., Puta, F. and Skruzny, M. 2004. Transcriptional coregulator SNW/SKIP: the concealed tie of dissimilar pathways. *Cell. Mol. Life Sci.* 61: 629-640.
6. Xu, C., Zhang, J., Huang, X., Sun, J., Xu, Y., Tang, Y., Wu, J., Shi, Y., Huang, Q. and Zhang, Q. 2006. Solution structure of human peptidyl prolyl isomerase-like protein 1 and insights into its interaction with SKIP. *J. Biol. Chem.* 281: 15900-15908.
7. Obama, K., Kato, T., Hasegawa, S., Satoh, S., Nakamura, Y. and Furukawa, Y. 2006. Overexpression of peptidyl-prolyl isomerase-like 1 is associated with the growth of colon cancer cells. *Clin. Cancer Res.* 12: 70-76.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: PPIL1 (human) mapping to 6p21.2.

PRODUCT

PPIL1 (h): 293T Lysate represents a lysate of human PPIL1 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

PPIL1 (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive PPIL1 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.