



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Dcp1a (h): 293T Lysate: sc-171459

BACKGROUND

Cleavage of the 5'-cap structure is involved in the major 5'-to-3' and non-sense-mediated mRNA decay pathways. The protein complex consisting of Dcp1 and Dcp2 has been identified as the species responsible for the decapping reaction in *Saccharomyces cerevisiae*. In nonsense-mediated decay, the human decapping complex, made up of *S. cerevisiae* homologs Dcp1a and hDcp2, may be recruited to mRNAs containing premature termination codons by nonsense-mediated decay factor (Upf) proteins. hDcp2 specifically hydrolyzes methylated capped RNA to release m⁷GDP, thereby aiding in mRNA degradation. Both Dcp1a and hDcp2 colocalize in the cytoplasm. In addition, Dcp1a interacts with Smad4 forming a complex with TGFβ and BMP-4. Dcp1a and Smad4 interact directly through a EVH1/WH1 domain on Dcp1a and a proline-rich activation domain on Smad4. Smad4 is essential to nuclear translocation of Dcp1a as deletion of the Smad4-interacting domain (located in the N-terminal 100 amino acids) of Dcp1a eliminates TGFβ-induced nuclear translocation of Dcp1a.

REFERENCES

1. toh, S., Itoh, F., Goumans, M.J. and Ten Dijke, P. 2000. Signaling of transforming growth factor-β family members through Smad proteins. *Eur. J. Biochem.* 267: 6954-6967.
2. Moustakas, A., Souchelnytskyi, S. and Heldin, C.H. 2001. Smad regulation in TGFβ signal transduction. *J. Cell Sci.* 114: 4359-4369.
3. Chen, W., Fu, X. and Sheng, Z. 2002. Review of current progress in the structure and function of Smad proteins. *Chin. Med. J.* 115: 446-450.
4. Bai, R.Y., Koester, C., Ouyang, T., Hahn, S.A., Hammerschmidt, M., Peschel, C. and Duyster, J. 2002. SMIF, a Smad4-interacting protein that functions as a co-activator in TGFβ signalling. *Nat. Cell. Biol.* 4: 181-190.
5. Callebaut, I. 2002. An EVH1/WH1 domain as a key actor in TGFβ signalling. *FEBS Lett.* 519: 178-180.

CHROMOSOMAL LOCATION

Genetic locus: DCP1A (human) mapping to 3p21.1.

PRODUCT

Dcp1a (h): 293T Lysate represents a lysate of human Dcp1a transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

Dcp1a (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive Dcp1a antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

Dcp1a (56-Y): sc-100706 is recommended as a positive control antibody for Western Blot analysis of enhanced human Dcp1a expression in Dcp1a transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



Dcp1a (56-Y): sc-100706. Western blot analysis of Dcp1a expression in non-transfected: sc-117752 (A) and human Dcp1a transfected: sc-171459 (B) 293T whole cell lysates.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.