



**SZABO  
SCANDIC**

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](http://linkedin.com/company/szaboscandic)



# CYFIP1 (h): 293T Lysate: sc-171687

## BACKGROUND

Fragile X syndrome is the most frequent form of inherited mental retardation and is a result of transcriptional silencing of the FMR1 gene on the X chromosome. The FMR1 protein (also designated FMRP) is an RNA-binding protein that associates with polyribosomes and is a likely component of a messenger ribonuclear protein (mRNP) particle. FMR1 can also interact with two fragile X syndrome related factors, FXR1 (also designated FXR1P) and FXR2 (also designated FXR2P). These proteins form heterodimers through their N-terminal coiled-coiled domains. CYFIP1 and CYFIP2 (also known as cytoplasmic FMRP interacting proteins 1 and 2, respectively, and as Sra-1 in mouse) both interact with FMR1 but CYFIP2 also reacts with FXR1 and FXR2. CYFIP1 and CYFIP2 bind GTP-bound Rac1 to release FMRP in its active state, which is thought to regulate mRNA translation of neural cytoskeletal proteins. A loss of CYFIP1 and CYFIP2 leads to mutant neurons with defective axonal growth and motor function.

## REFERENCES

1. Schenck, A., Bardoni, B., Moro, A., Bagni, C. and Mandel, J.L. 2001. A highly conserved protein family interacting with the fragile X mental retardation protein (FMRP) and displaying selective interactions with FMRP-related proteins FXR1P and FXR2P. *Proc. Natl. Acad. Sci. USA* 98: 8844-8849.
2. Kunda, P., Craig, G., Dominguez, V. and Baum, B. 2003. Abi, Sra-1, and Kette control the stability and localization of SCAR/WAVE to regulate the formation of Actin-based protrusions. *Curr. Biol.* 13: 1867-1875.
3. Billuart, P. and Chelly, J. 2003. From fragile X mental retardation protein to Rac1 GTPase: new insights from Fly CYFIP. *Neuron* 38: 843-845.
4. Schenck, A., Bardoni, B., Langmann, C., Harden, N., Mandel, J.L. and Giangrande, A. 2003. CYFIP/Sra-1 controls neuronal connectivity in *Drosophila* and links the Rac1 GTPase pathway to the fragile X protein. *Neuron* 38: 887-898.
5. Schenck, A., Qurashi, A., Carrera, P., Bardoni, B., Diebold, C., Schejter, E., Mandel, J.L. and Giangrande, A. 2004. WAVE/SCAR, a multifunctional complex coordinating different aspects of neuronal connectivity. *Dev. Biol.* 274: 260-270.
6. Brembu, T., Winge, P., Seem, M. and Bones, A.M. 2004. NAPP and PIRP encode subunits of a putative wave regulatory protein complex involved in plant cell morphogenesis. *Plant Cell* 16: 2335-2349.
7. Bogdan, S., Grewe, O., Strunk, M., Mertens, A. and Klamt, C. 2004. Sra-1 interacts with Kette and Wasp and is required for neuronal and bristle development in *Drosophila*. *Development* 131: 3981-3989.
8. Handa, V., Goldwater, D., Stiles, D., Cam, M., Poy, G., Kumari, D. and Usdin, K. 2005. Long CGG-repeat tracts are toxic to human cells: implications for carriers of fragile X premutation alleles. *FEBS Lett.* 579: 2702-2708.
9. Pilpel, Y. and Segal, M. 2005. Rapid WAVE dynamics in dendritic spines of cultured hippocampal neurons is mediated by Actin polymerization. *J. Neurochem.* 95: 1401-1410.

## CHROMOSOMAL LOCATION

Genetic locus: CYFIP1 (human) mapping to 15q11.2.

## PRODUCT

CYFIP1 (h): 293T Lysate represents a lysate of human CYFIP1 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

## APPLICATIONS

CYFIP1 (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive CYFIP1 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

## STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.