



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

ISYNA1 (h2): 293T Lysate: sc-171752

BACKGROUND

Myo-inositol is an important constituent of membrane phospholipids and is a precursor for the phosphoinositide signaling pathway. ISYNA1 (inositol-3-phosphate synthase 1), also known as IPS, INO1 or INOS, is a 558 amino acid enzyme belonging to the myo-inositol-1-phosphate synthase family. Highly expressed in testis, ovary, heart, placenta and pancreas, with weak expression in blood leukocytes, thymus, skeletal muscle and colon, SYNA1 is the key enzyme myo-inositol biosynthesis, as it catalyzes the conversion of glucose 6-phosphate to 1-myoinositol 1-phosphate in a NAD-dependent manner. ISYNA1 is the rate-limiting enzyme in the synthesis of all inositol-containing compounds. ISYNA1 may be upregulated by E2F-1, and is inhibited by valproate (VPA) and lithium, which are mood-stabilizing drugs.

REFERENCES

- Hegeman, C.E., Good, L.L. and Grabau, E.A. 2001. Expression of D-myoinositol-3-phosphate synthase in soybean. Implications for phytic acid biosynthesis. *Plant Physiol.* 125: 1941-1948.
- Agam, G., Shamir, A., Shaltiel, G. and Greenberg, M.L. 2002. Myo-inositol-1-phosphate (MIP) synthase: a possible new target for antibipolar drugs. *Bipolar Disord.* 4: 15-20.
- Seelan, R.S., Parthasarathy, L.K. and Parthasarathy, R.N. 2004. E2F1 regulation of the human myo-inositol 1-phosphate synthase (ISYNA1) gene promoter. *Arch. Biochem. Biophys.* 43: 95-106.
- Chauvin, T.R. and Griswold, M.D. 2004. Characterization of the expression and regulation of genes necessary for myo-inositol biosynthesis and transport in the seminiferous epithelium. *Biol. Reprod.* 70: 744-751.
- Ju, S., Shaltiel, G., Shamir, A., Agam, G. and Greenberg, M.L. 2004. Human 1-D-myoinositol-3-phosphate synthase is functional in yeast. *J. Biol. Chem.* 279: 21759-21765.
- Groenen, P.M., Klootwijk, R., Schijvenaars, M.M., Straatman, H., Mariman, E.C., Franke, B. and Steegers-Theunissen, R.P. 2004. Spina bifida and genetic factors related to myo-inositol, glucose, and zinc. *Mol. Genet. Metab.* 82: 154-161.
- Shaltiel, G., Galit, S., Mark, S., Shirley, M., Kofman, O., Ora, K., Belmaker, R.H., Agam, G. and Galila, A. 2007. Effect of valproate derivatives on human brain myo-inositol-1-phosphate (MIP) synthase activity and amphetamine-induced rearing. *Pharmacol. Rep.* 59: 402-407.
- Einat, H., Tian, F., Belmaker, R.H. and Frost, J.W. 2008. Myo-inositol-1-phosphate (MIP) synthase inhibition: *in vivo* study in rats. *J. Neural. Transm.* 115: 55-58.
- Seelan, R.S., Lakshmanan, J., Casanova, M.F. and Parthasarathy, R.N. 2009. Identification of myo-inositol-3-phosphate synthase isoforms: characterization, expression, and putative role of a 16-kDa γ_c isoform. *J. Biol. Chem.* 284: 9443-9457.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

CHROMOSOMAL LOCATION

Genetic locus: ISYNA1 (human) mapping to 19p13.11.

PRODUCT

ISYNA1 (h2): 293T Lysate represents a lysate of human ISYNA1 transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

ISYNA1 (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive ISYNA1 antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.