



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

FBP1 (h2): 293T Lysate: sc-171763

BACKGROUND

Activation of FUSE, the far-upstream element, is required for the proper expression of the mammalian gene c-Myc in undifferentiated cells. The binding of FBP (FUSE-binding protein or far-upstream element binding protein) to FUSE is necessary for c-Myc expression, indicating that FBP functions as a growth-dependent regulator of c-Myc expression. Isolated from proliferating HL-60 cells, FBP, FBP2 and FBP3 comprise a family of single-stranded DNA-binding proteins that specifically bind to FUSE elements. The FBP transcription factors share a conserved central DNA-binding domain and show significant homology in their carboxyl-terminal activation domains. Expression of FBP is detected in undifferentiated cells and is substantially decreased following cellular differentiation.

REFERENCES

1. Avigan, M.I., et al. 1990. A far-upstream element stimulates c-Myc expression in undifferentiated leukemia cells. *J. Biol. Chem.* 265: 18538-18545.
2. Duncan, R.D., et al. 1994. A sequence-specific, single strand binding protein activates the far-upstream of c-Myc and defines a new DNA-binding motif. *Genes Dev.* 8: 465-480.
3. Bazar, L., et al. 1995. A transactivator of c-Myc is coordinately regulated with the proto-oncogene during cellular growth. *Oncogene* 10: 2229-2238.
4. Davis-Smyth, T., et al. 1996. The far-upstream element-binding proteins comprise an ancient family of single-strand DNA-binding transactivators. *J. Biol. Chem.* 271: 31679-31687.
5. Michelotti, G.A., et al. 1996. Multiple single-stranded *cis* elements are associated with activated chromatin of the human c-Myc gene *in vivo*. *Mol. Cell. Biol.* 16: 2656-2669.
6. Rehbein, M., et al. 2002. Molecular characterization of MARTA1, a protein interacting with the dendritic targeting element of MAP2 mRNAs. *J. Neurochem.* 82: 1039-1046.
7. Braddock, D.T., et al. 2002. Structure and dynamics of KH domains from FBP bound to single-stranded DNA. *Nature* 415:1051-1056.

CHROMOSOMAL LOCATION

Genetic locus: FUBP1 (human) mapping to 1p31.1.

PRODUCT

FBP1 (h2): 293T Lysate represents a lysate of human FBP1 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

APPLICATIONS

FBP1 (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive FBP1 antibodies.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.