



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



YSK1 (h3): 293T Lysate: sc-171799

BACKGROUND

Several mammalian kinases have been identified which exhibit sequence similarity to the *Saccharomyces cerevisiae* serine/threonine kinase Ste20. Ste20 is involved in relaying signals from G protein-coupled receptors, and it lies upstream of a MAP kinase kinase kinase. Mammalian Ste20-like kinases include YSK1, KHS, GLK, NIK, HPK1, Krs-1, Krs-2 and GC kinase. YSK1 (yeast SPS/Ste20-related kinase 1) is expressed in a wide variety of cell types and tissues and has been shown to have kinase activity. Unlike many of the other Ste20-like kinases, however, overexpression of YSK1 does not lead to activation of the SAPK/JNK pathway.

REFERENCES

1. Leberer, E., Dignard, D., Harcus, D., Thomas, D.Y. and Whiteway, M. 1992. The protein kinase homologue Ste20p is required to link the yeast pheromone response G protein $\beta\gamma$ subunits to downstream signalling components. EMBO J. 11: 4815-4824.
2. Wu, C., Whiteway, M., Thomas, D.Y. and Leberer, E. 1995. Molecular characterization of Ste20p, a potential mitogen-activated protein or extracellular signal-regulated kinase kinase (MEK) kinase kinase from *Saccharomyces cerevisiae*. J. Biol. Chem. 270: 15984-15992.
3. Su., Y.C., Han, J., Xu, S., Cobb, M. and Skolnik, E.Y. 1997. NIK is a new Ste20-related kinase that binds NCK and MEKK1 and activates the SAPK/JNK cascade via a conserved regulatory domain. EMBO J. 16: 1279-1290.
4. Diener, K., Wang, X.S., Chen, C., Meyer, C.F., Keesler, G., Zukowski, M., Tan, T.H. and Yao, Z. 1997. Activation of the c-Jun N-terminal kinase pathway by a novel protein kinase related to human germinal center kinase. Proc. Natl. Acad. Sci. USA 94: 9687-9692.
5. Osada, S., Izawa, M., Saito, R., Mizuno, K., Suzuki, A., Hirai, S. and Ohno, S. 1997. YSK1, a novel mammalian protein kinase structurally related to Ste20 and SPS1, but is not involved in the known MAPK pathways. Oncogene 14: 2047-2057.

CHROMOSOMAL LOCATION

Genetic locus: STK25 (human) mapping to 2q37.3.

PRODUCT

YSK1 (h3): 293T Lysate represents a lysate of human YSK1 transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

YSK1 (h3): 293T Lysate is suitable as a Western Blotting positive control for human reactive YSK1 antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.