



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Exo1 (h2): 293T Lysate: sc-172184

BACKGROUND

Comparative evaluation of the expression patterns of the human and mouse genes, combined with previous biochemical and yeast genetic studies, indicate that the Exo1 (Exonuclease I) proteins are important contributors to chromosome processing during mammalian DNA repair and recombination. In mice, the Exo1 gene maps to distal chromosome 1, consistent with the recent mapping of the orthologous human HEX1/EXO1 gene to chromosome 1q43. Exo1 is expressed prominently in testis, an area of active homologous recombination, and spleen, a prominent lymphoid tissue. In both mammalian and yeast systems, Exo1 is a 5'-3' double stranded DNA exonuclease that has previously been implicated in DNA mismatch repair (MMR). The MMR system ensures genome integrity by removing mispaired and unpaired bases that originate during replication. In humans, Exo1 interacts with MSH2 and MLH1 and has been proposed to be a redundant exonuclease in MMR. In both mammalian and yeast systems, Exo1 plays a structural role in MMR and stabilizes multi-protein complexes containing a number of MMR proteins.

REFERENCES

1. Lee, B.I., Shannon, M., Stubbs, L. and Wilson, D.M. III 1999. Expression specificity of the mouse exonuclease 1 (mExo1) gene. *Nucleic Acids Res.* 27: 4114-4120.
2. Kirkpatrick, D.T., Ferguson, J.R., Petes, T.D. and Symington, L.S. 2000. Decreased meiotic intergenic recombination and increased meiosis I non-disjunction in Exo1 mutants of *Saccharomyces cerevisiae*. *Genetics* 156: 1549-1557.
3. Tran, P.T., Simon, J.A. and Liskay, R.M. 2001. Interactions of Exo1p with components of MutL α in *Saccharomyces cerevisiae*. *Proc. Natl. Acad. Sci. USA* 98: 9760-9765.
4. Mansour, A.A., Tornier, C., Lehmann, E., Darmon, M. and Fleck, O. 2001. Control of GT repeat stability in *Schizosaccharomyces pombe* by mismatch repair factors. *Genetics* 158: 77-85.
5. Amin, N.S., Nguyen, M.N., Oh, S. and Kolodner, R.D. 2001. Exo1-dependent mutator mutations: model system for studying functional interactions in mismatch repair. *Mol. Cell. Biol.* 21: 5142-5155.
6. Tran, P.T., Erdeniz, N., Symington, L.S. and Liskay, R.M. 2004. EXO1-A multi-tasking eukaryotic nuclease. *DNA repair.* 3: 1549-59.
7. Barber, L.J., Ward, T.A., Hartley, J.A. and McHugh, P.J. 2005. DNA inter-strand cross-link repair in the *Saccharomyces cerevisiae* cell cycle: overlapping roles for PSO2 (Snm1) with MutS factors and Exo1 during S phase. *Mol. Cell. Biol.* 25: 2297-2309.
8. Cotta-Ramusino, C., Fachinetti, D., Lucca, C., Doksani, Y., Lopes, M., Sogo, J. and Foiani, M. 2005. Exo1 processes stalled replication forks and counteracts fork reversal in checkpoint-defective cells. *Mol. Cell* 17: 153-159.
9. Yamamoto, H., Hanafusa, H., Ouchida, M., Yano, M., Suzuki, H., Murakami, M., Aoe, M., Shimizu, N., Nakachi, K. and Shimizu, K. 2005. Single nucleotide polymorphisms in the EXO1 gene and risk of colorectal cancer in a Japanese population. *Carcinogenesis* 26: 411-416.

CHROMOSOMAL LOCATION

Genetic locus: EXO1 (human) mapping to 1q43.

PRODUCT

Exo1 (h2): 293T Lysate represents a lysate of human Exo1 transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

Exo1 (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive Exo1 antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.