



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

ITM2B (h): 293T Lysate: sc-172355

BACKGROUND

The type II integral membrane (ITM2) protein family consists of three members: ITM2A (also designated E25), ITM2B and ITM2C. ITM2A expression is high in osteogenic and lymphoid tissues, while both ITM2B and ITM2C are expressed in brain. ITM2B is a 266 amino acid protein that contains a potential N-glycosylation site, a potential single transmembrane-spanning domain between amino acids 52 and 74, and an extracellular C-terminal domain. Mutations in the ITM2B gene can lead to familial British dementia (FBD), an autosomal dominant disease with an onset around the fifth decade of life that is characterized by progressive dementia, spasticity and cerebellar ataxia. Familial Danish dementia (FDD), also designated hereditary ophthalmoto-encephalica, is also associated with mutations in the ITM2B gene. FDD is an autosomal dominant disorder characterized by cataracts, deafness, progressive ataxia and dementia.

REFERENCES

- Ernst, B., Dalby, M.A. and Dalby, A. 1970. Luria testing in demented patients. *Acta Neurol. Scand.* 46: 97-98.
- Vidal, R., Frangione, B., Rostagno, A., Mead, S., Révész, T., Plant, G. and Ghiso, J. 1999. A stop-codon mutation in the BRI gene associated with familial British dementia. *Nature* 399: 776-781.
- Ghiso, J.A., Holton, J., Miravalle, L., Calero, M., Lashley, T., Vidal, R., Houlden, H., Wood, N., Neubert, T.A., Rostagno, A., Plant, G., Revesz, T. and Frangione, B. 2001. Systemic amyloid deposits in familial British dementia. *J. Biol. Chem.* 276: 43909-43914.
- El-Agnaf, O., Gibson, G., Lee, M., Wright, A. and Austen, B.M. 2004. Properties of neurotoxic peptides related to the Bri gene. *Protein Pept. Lett.* 11: 207-212.
- Matsuda, S., Giliberto, L., Matsuda, Y., Davies, P., McGowan, E., Pickford, F., Ghiso, J., Frangione, B. and D'Adamo, L. 2005. The familial dementia BRI2 gene binds the Alzheimer gene β -Amyloid precursor protein and inhibits β -Amyloid production. *J. Biol. Chem.* 280: 28912-28916.
- Zirn, B., Samans, B., Spangenberg, C., Graf, N., Eilers, M. and Gessler, M. 2005. All-*trans* retinoic acid treatment of Wilms tumor cells reverses expression of genes associated with high risk and relapse *in vivo*. *Oncogene* 24: 5246-5251.
- Morelli, L., Llovera, R.E., Alonso, L.G., Frangione, B., de Prat-Gay, G., Ghiso, J. and Castaño, E.M. 2005. Insulin-degrading enzyme degrades amyloid peptides associated with British and Danish familial dementia. *Biochem. Biophys. Res. Commun.* 332: 808-816.

CHROMOSOMAL LOCATION

Genetic locus: ITM2B (human) mapping to 13q14.2.

PRODUCT

ITM2B (h): 293T Lysate represents a lysate of human ITM2B transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

ITM2B (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive ITM2B antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.