



**SZABO  
SCANDIC**

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](http://linkedin.com/company/szaboscandic)



# IRX3 (h): 293T Lysate: sc-173075

## BACKGROUND

The Iroquois homeobox gene family of transcription factors regulate aspects of embryonic development including anterior/posterior and dorsal/ventral axis patterning in the central nervous system. The Iroquois family are clustered on two loci, IRXA and IRXB, which map to chromosomes 8 and 13 in mice. The IRXA group includes IRX1, IRX2 and IRX4; the IRXB group comprises IRX3, IRX5 and IRX6. IRX1 and IRX2 are both widely expressed during development in the lung epithelium and also in the ventricular septum. IRX1 and IRX2 also play a role in digit formation (E11.5-E14.5). The IRX gene family members are each expressed in a distinct pattern during mouse heart development. Specifically, IRX1 and IRX2 are expressed in the ventricular septum and IRX3 is expressed in the ventricular trabeculated myocardium. In addition, IRX4 is expressed in the linear heart tube and the AV canal; IRX5 is expressed in the endocardium lining the ventricular and atrial myocardium. Furthermore, the IRX4 gene may modulate cardiac development and function. Although the heart of IRX4<sup>-/-</sup> mice appears to develop normally, adult IRX4<sup>-/-</sup> mice exhibit cardiomyopathy, including cardiac hypertrophy and decreased contractility.

## REFERENCES

- Christoffels, V.M., et al. 2000. Patterning the embryonic heart: identification of five mouse Iroquois homeobox genes in the developing heart. *Dev. Biol.* 224: 263-274.
- Mummendorff, J., et al. 2001. Expression of IRX6 during mouse morphogenesis. *Mech. Dev.* 103: 193-195.
- Becker, M.B., et al. 2001. IRX1 and IRX2 expression in early lung development. *Mech. Dev.* 106: 155-158.
- Zulch, A., et al. 2001. Expression pattern of IRX1 and IRX2 during mouse digit development. *Mech. Dev.* 106: 159-162.
- Bruneau, B.G., et al. 2001. Cardiomyopathy in IRX4-deficient mice is preceded by abnormal ventricular gene expression. *Mol. Cell. Biol.* 21: 1730-1736.
- Ogura, K., et al. 2001. Cloning and chromosome mapping of human and chicken Iroquois (IRX) genes. *Cytogenet. Cell Genet.* 92: 320-325.
- Online Mendelian Inheritance in Man, OMIM™. 2001. Johns Hopkins University, Baltimore, MD. MIM Number: 606197. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

## CHROMOSOMAL LOCATION

Genetic locus: IRX3 (human) mapping to 16q12.2.

## PRODUCT

IRX3 (h): 293T Lysate represents a lysate of human IRX3 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

## STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## APPLICATIONS

IRX3 (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive IRX3 antibodies.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.


IRX3 (G-6): sc-166877 is recommended as a positive control antibody for Western Blot analysis of enhanced human IRX3 expression in IRX3 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:

1) Western Blotting: use m-IgG<sub>κ</sub> BP-HRP: sc-516102 or m-IgG<sub>κ</sub> BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

## DATA



IRX3 (G-6): sc-166877. Western blot analysis of IRX3 expression in non-transfected: sc-117752 (**A**) and human IRX3 transfected: sc-173075 (**B**) 293T whole cell lysates.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.