



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



DR4 (h2): 293T Lysate: sc-173513

BACKGROUND

Tumor necrosis factor (TNF) is a pleiotropic cytokine whose function is mediated by two distinct cell surface receptors, designated TNF-R1 and TNF-R2, which are expressed on most cell types. TNF function is primarily mediated through TNF-R1 signaling. Both receptors belong to the growing TNF receptor superfamily which includes FAS antigen and CD40. TNF-R1 contains a cytoplasmic motif, termed the "death domain," that has been found to be necessary for the transduction of the apoptotic signal. The death domain is also found in several other receptors, including FAS, DR2 (or TRUNDD), DR3 (death receptor 3), DR4 and DR5. TRUNDD, DR4 and DR5 are receptors for the apoptosis-inducing cytokine TRAIL. A non-death domain-containing receptor, designated decoy receptor (DcR1 or TRID), also specifically associates with TRAIL and may play a role in cellular resistance to apoptotic stimuli.

REFERENCES

1. Tartaglia, L.A., Ayres, T.M., Wong, G.H. and Goeddel, D.V. 1993. A novel domain within the 55 kd TNF receptor signals cell death. *Cell* 74: 845-853.
2. Smith, C.A., Farrah, T. and Goodwin, R.G. 1994. The TNF receptor superfamily of cellular and viral proteins: activation, costimulation, and death. *Cell* 76: 959-962.
3. Nagata, S. and Golstein, P. 1995. The FAS death factor. *Science* 267: 1449-1456.
4. Pan, G., Ni, J., Yu, G., Wei, Y.F. and Dixit, V.M. 1998. TRUNDD, a new member of the TRAIL receptor family that antagonizes TRAIL signalling. *FEBS Lett.* 424: 41-45.
5. Kitson, J., Raven, T., Jiang, Y.P., Goeddel, D.V., Giles, K.M., Pun, K.T., Grinham, C.J., Brown, R. and Farrow, S.N. 1996. A death-domain-containing receptor that mediates apoptosis. *Nature* 384: 372-375.
6. Pan, G., O'Rourke, K., Chinnaiyan, A.M., Gentz, R., Ebner, R., Ni, J. and Dixit, V.M. 1997. The receptor for the cytotoxic ligand TRAIL. *Science* 276: 111-113.
7. Pan, G., Ni, J., Wei, Y.F., Yu, G., Gentz, R. and Dixit, V.M. 1997. An antagonist decoy receptor and a death domain-containing receptor for TRAIL. *Science* 277: 815-818.
8. Sheridan, J.P., Marsters, S.A., Pitti, R.M., Gurney, A., Skubatch, M., Baldwin, D., Ramakrishnan, L., Gray, C.L., Baker, K., Wood, W.I., Goddard, A.D., Godowski, P. and Ashkenazi, A. 1997. Control of TRAIL-induced apoptosis by a family of signaling and decoy receptors. *Science* 277: 818-821.

CHROMOSOMAL LOCATION

Genetic locus: TNFRSF10A (human) mapping to 8p21.3.

PRODUCT

DR4 (h2): 293T Lysate represents a lysate of human DR4 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.


APPLICATIONS

DR4 (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive DR4 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

DR4 (B-N28): sc-65312 is recommended as a positive control antibody for Western Blot analysis of enhanced human DR4 expression in DR4 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

DATA



DR4 (B-N28): sc-65312. Western blot analysis of DR4 expression in non-transfected: sc-117752 (**A**) and human DR4 transfected: sc-173513 (**B**) 293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.