



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TRIM8 (h): 293T Lysate: sc-174016

BACKGROUND

The tripartite motif (TRIM) family of proteins are characterized by a conserved TRIM domain that includes a coiled-coil region, a B-box type zinc finger, one RING finger and three zinc-binding domains. TRIM8 (tripartite motif containing 8), also known as GERP (glioblastoma-expressed RING finger protein) or RNF27 (RING finger protein 27), is a 551 amino acid protein that is thought to function as an E3 ubiquitin-protein ligase that promotes SOCS-1 proteasomal degradation. As a widely expressed homodimer, TRIM8 localizes to nuclear bodies and contains two B box-type zinc fingers and one RING-type zinc finger. TRIM8 is expressed in lung, heart, brain and skeletal muscle, with low levels detected in intestine, placenta, leukocytes and liver. The gene encoding TRIM8 maps to human chromosome 10q24.32.

REFERENCES

1. Vincent, S.R., Kwasnicka, D.A. and Fretier, P. 2000. A novel RING finger-B box-coiled-coil protein, GERP. *Biochem. Biophys. Res. Commun.* 279: 482-486.
2. Reymond, A., Meroni, G., Fantozzi, A., Merla, G., Cairo, S., Luzi, L., Riganelli, D., Zanaria, E., Messali, S., Cainarca, S., Guffanti, A., Minucci, S., Pelicci, P.G. and Ballabio, A. 2001. The tripartite motif family identifies cell compartments. *EMBO J.* 20: 2140-2151.
3. Toniato, E., Chen, X.P., Losman, J., Flati, V., Donahue, L. and Rothman, P. 2002. TRIM8/GERP RING finger protein interacts with SOCS-1. *J. Biol. Chem.* 277: 37315-37322.
4. Toniato, E., Flati, V., Laglia, E., Mincione, G. and Martinotti, S. 2004. Genomic organization and cytokine-mediated inducibility of the human TRIM-8/Gerp gene. *Int. J. Immunopathol. Pharmacol.* 17: 71-76.
5. Online Mendelian Inheritance in Man, OMIM™. 2007. Johns Hopkins University, Baltimore, MD. MIM Number: 606125. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
6. Markson, G., Kiel, C., Hyde, R., Brown, S., Charalabous, P., Bremm, A., Semple, J., Woodsmith, J., Duley, S., Salehi-Ashtiani, K., Vidal, M., Komander, D., Serrano, L., Lehner, P. and Sanderson, C.M. 2009. Analysis of the human E2 ubiquitin conjugating enzyme protein interaction network. *Genome Res.* 19: 1905-1911.
7. Okumura, F., Matsunaga, Y., Katayama, Y., Nakayama, K.I. and Hatakeyama, S. 2010. TRIM8 modulates STAT3 activity through negative regulation of PIAS3. *J. Cell Sci.* 123: 2238-2245.

CHROMOSOMAL LOCATION

Genetic locus: TRIM8 (human) mapping to 10q24.32.

PRODUCT

TRIM8 (h): 293T Lysate represents a lysate of human TRIM8 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

TRIM8 (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive TRIM8 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.