



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Rho 8 (h): 293T Lysate: sc-174030

BACKGROUND

The Ras p21 family of guanine nucleotide proteins has been widely studied in view of its apparent role in signal transduction pathways and high frequency of mutations in human malignancies. It is now clear, however, that the Ras proteins (H-, K- and N-Ras p21) are members of a much larger superfamily of related proteins. Six members of this family, Rap 1A, Rap 1B, Rap 2, R-Ras, Ral A and Ral B, exhibit approximately 50% amino acid homology to Ras. The six mammalian Rho proteins (Rho A, B, C, G, 7 and 8) are approximately 30% homologous to Ras and are expressed in a wide range of cell types. Both Ras p21 and Rho p21, as well as other members of the Ras superfamily, contain a carboxy-terminal CAAX sequence (C, cysteine; A, aliphatic amino acid; X, any amino acid) which in the case of Ras has been shown to be essential for correct localization and function.

REFERENCES

1. Madaule, P. and Axel R. 1985. A novel Ras-related gene family. *Cell* 41: 31-40.
2. Yeramian, P., Chardin, P., Madaule, P. and Tavitian, A. 1987. Nucleotide sequence of human Rho cDNA clone 12. *Nucleic Acids Res.* 15: 1869.
3. Barbacid, M. 1987. Ras genes. *Annu. Rev. Biochem.* 56: 779-827.
4. Olofsson, B., Chardin, P., Touchot, N., Zahraoui, A. and Tavitian, A. 1988. Expression of the Ras-related Ral A Rho 12 and Rab genes in adult mouse tissues. *Oncogene* 3: 231-234.
5. Chardin, P. 1988. The Ras superfamily proteins. *Biochimie* 70: 865-868.
6. Morris, J.D., et al. 1989. Scrape-loading of Swiss 3T3 cells with Ras protein rapidly activates protein kinase C in the absence of phosphoinositide hydrolysis. *Oncogene* 4: 27-31.
7. Garrett, M.D., Price, B., Lloyd, A.C., Self, A.J., Marshall, C.J. and Hall, A. 1989. Identification of distinct cytoplasmic targets for Ras/R-Ras and Rho regulatory proteins. *J. Biol. Chem.* 264: 10-13.
8. Adamson, P., Marshall, C.J., Hall, A. and Tilbrook, P.A. 1992. Post-translational modifications of p21 Rho proteins. *J. Biol. Chem.* 267: 20033-20038.

CHROMOSOMAL LOCATION

Genetic locus: RND3 (human) mapping to 2q23.3.

PRODUCT

Rho 8 (h): 293T Lysate represents a lysate of human Rho 8 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

APPLICATIONS

Rho 8 (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive Rho 8 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.