



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

SEN8 (h2): 293T Lysate: sc-174054

BACKGROUND

SUMO (small ubiquitin-related modifier), a member of the ubiquitin-like protein family, regulates diverse cellular functions of a variety of target proteins, including transcription, DNA repair, nucleocytoplasmic trafficking and chromosome segregation. SUMO precursor proteins undergo cleavage of the residues after the "GG" region by SUMO-specific proteases in maturation. This cleavage of the precursor is a prerequisite for subsequent sumoylation. The sentrin-specific (or SUMO-specific) protease (SEN) proteins belong to the peptidase C48 family and include SEN1-3 and SEN5-8. SEN1, SEN2 and SEN3 degrade UBL1 and SMT3H2 conjugates and subsequently release the monomers from sumoylated substrates. HIPK2 is a desumoylation target for SEN1 which shuttles between the cytoplasm and the nucleus. Mutation analyses reveal that SEN1 contains the nuclear export sequence (NES) within the extreme carboxyl-terminal region, and SEN1 is exported to the cytoplasm in a NES-dependent manner. SEN2 has been implicated as a downregulator of CTNNB1 levels and may therefore be a modulator of the Wnt pathway. SUMO protease SEN3 reverses the sumoylation of MEF2 to augment its transcriptional and myogenic activities. SEN5 localizes to the nucleolus and preferentially processes SUMO-3. It is thought to play a role in mitosis and/or cytokinesis. SEN6 localizes to the cytoplasm and releases SUMO-1. Expression of SEN6 is higher in reproductive organs, indicating that it may mediate processes related to reproduction. SEN8 is involved in the release of sentrins.

REFERENCES

1. Gong, L., et al. 2000. Differential regulation of sentrinized proteins by a novel sentrin-specific protease. *J. Biol. Chem.* 275: 3355-3359.
2. Kim, K.I., et al. 2000. A new SUMO-1-specific protease, SUSP1, that is highly expressed in reproductive organs. *J. Biol. Chem.* 275: 14102-14106.
3. Cheng, J., et al. 2004. SEN1 enhances androgen receptor-dependent transcription through desumoylation of histone deacetylase 1. *Mol. Cell. Biol.* 24: 6021-6028.
4. Reverter, D., et al. 2004. A basis for SUMO protease specificity provided by analysis of human SEN2 and a SEN2-SUMO complex. *Structure* 12: 1519-1531.
5. Kim, Y.H., et al. 2005. Desumoylation of homeodomain-interacting protein kinase 2 (HIPK2) through the cytoplasmic-nuclear shuttling of the SUMO-specific protease SEN1. *FEBS Lett.* 579: 6272-6278.
6. Xu, Z., et al. 2005. Mapping residues of SUMO precursors essential in differential maturation by SUMO-specific protease, SEN1. *Biochem. J.* 386: 325-330.
7. Gong, L. and Yeh, E.T. 2006. Characterization of a family of nucleolar SUMO-specific proteases with preference for SUMO-2 or SUMO-3. *J. Biol. Chem.* 281: 15869-15877.
8. Di Bacco, A., et al. 2006. The SUMO-specific protease SEN5 is required for cell division. *Mol. Cell. Biol.* 26: 4489-4498.
9. Shen, L.N., et al. 2006. The structure of SEN1- between SUMO paralogues during processing. *Biochem. J.* 397: 279-288.

CHROMOSOMAL LOCATION

Genetic locus: SEN8 (human) mapping to 15q23.

PRODUCT

SEN8 (h2): 293T Lysate represents a lysate of human SEN8 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

SEN8 (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive SEN8 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.