



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Calpain 10 (h): 293T Lysate: sc-174154

BACKGROUND

The CAPN10 (Calpain 10) gene encodes a ubiquitously expressed member of the Calpain-like cysteine protease family and shows association with type 2 diabetes research suggests that Calpain 10 plays a role in an innovative pathway involved in the pathophysiology of diabetes, where Calpain-10 represents the third example of a protease contributing to the advancement of diabetes, the others being prohormone convertase-1 and prohormone-processing carboxypeptidase E, both of which are associated with diabetes and obesity. The CAPN10 human cDNA encodes a 672 amino-acid protein that shares 81.7% identity with the mouse ortholog, and analysis of human cDNA clones displays an intricate pattern of alternative splicing. CAPN10, which presumably plays a role in the regulation of Insulin secretion, is thought to contain a signature of the effects of positive natural selection within its genetic sequence.

REFERENCES

1. Horikawa, Y., Oda, N., Cox, N.J., Li, X., Orho-Melander, M., Hara, M., Hinokio, Y., Lindner, T.H., Mashima, H., Schwarz, P.E., del Bosque-Plata, L., Horikawa, Y., Oda, Y., Yoshiuchi, I., Colilla, S., Polonsky, K.S., Wei, S., Concannon, P., Iwasaki, N., Schulze, J., Baier, L.J., Bogardus, C., Groop, L., et al. 2000. Genetic variation in the gene encoding Calpain-10 is diabetes mellitus. *Nat. Genet.* 26: 163-175.
2. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 605286. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
3. Clark, V.J., Cox, N.J., Hammond, M., Hanis, C.L. and Di Rienzo, A. 2005. Haplotype structure and phylogenetic shadowing of a hypervariable region in the CAPN10 gene. *Nat. Genet.* 117: 258-266.
4. Ridderstrale, M., Parikh, H. and Groop, L. 2005. Calpain 10 and type 2 diabetes: are we getting closer to an explanation? *Curr. Opin. Clin. Nutr. Metab. Care* 8: 361-366.
5. Vander Molen, J., Frisse, L.M., Fullerton, S.M., Qian, Y., Del Bosque-Plata, L., Hudson, R.R. and Di Rienzo, A. 2005. Population genetics of CAPN10 and GPR35: implications for the evolution of type 2 diabetes variants. *Am. J. Hum. Genet.* 76: 548-560.
6. Wolford, J.K. and Vozarova de Courten, B. 2005. Genetic basis of type 2 diabetes mellitus: implications for therapy. *Treat. Endocrinol.* 3: 257-267.
7. Wu, B., Takahashi, J., Fu, M., Cheng, H., Matsumura, S. and Taniguchi, H. 2005. Variants of Calpain-10 gene and its association with type 2 diabetes mellitus in a Chinese population. *Diabetes Res. Clin. Pract.* 68: 155-161.
8. Tsuchiya, T., Schwarz, P.E., Bosque-Plata, L.D., Geoffrey Hayes, M., Dina, C., Froguel, P., Wayne Towers, G., Fischer, S., Temelkova-Kurktschiev, T., Rietzsch, H., Graessler, J., Vcelák, J., Palyzová, D., Selisko, T., Bendlová, B., Schulze, J., Julius, U., Hanefeld, M., Weedon, M.N., Evans, J.C., et al. 2006. Association of the Calpain-10 gene with type 2 diabetes in Europeans: of pooled and meta-analyses. *Mol. Genet. Metab.* 89: 174-184.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

CHROMOSOMAL LOCATION

Genetic locus: CAPN10 (human) mapping to 2q37.3.

PRODUCT

Calpain 10 (h): 293T Lysate represents a lysate of human Calpain 10 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

Calpain 10 (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive Calpain 10 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.