



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

OGG1/2 (h): 293T Lysate: sc-174305

BACKGROUND

8-oxoguanine (8-oxoG), an oxidized form of guanine, is produced by reactive oxygen species in both DNA and nucleotide pools during normal aging. Accumulation of 8-oxoG increases the occurrence of A:T to C:G or G:C to T:A transversion mutations, because 8-oxoG forms a stable basepair with adenine as well as with cytosine. OGG1 (for 8-oxoG DNA glycosylase), also designated MMH, is a DNA repair enzyme that corrects these mutations. Inactivation of the OGG1 gene leads to a mutator phenotype, characterized by the increase in G:C to T:A transversions. The OGG1 gene encodes eight isoforms (OGG1A-C, OGG2A-E) which result from alternative splicing of a single messenger RNA. The OGG1A splice variant is the most prevalent form and localizes to the nucleus, whereas the OGG2A splice variant is targeted to the mitochondria.

REFERENCES

1. Shibutani, S., Takeshita, M. and Grollman, A.P. 1991. Insertion of specific bases during DNA synthesis past the oxidation-damaged base 8-oxodG. *Nature* 349: 431-434.
2. Cheng, K.C., Cahill, D.S., Kasai, H., Nishimura, S. and Loeb, L.A. 1992. 8-hydroxyguanine, an abundant form of oxidative DNA damage, causes GT and AC substitutions. *J. Biol. Chem.* 267: 166-172.
3. Ames, B.N., Shigenaga, M.K. and Hagen, T.M. 1993. Oxidants, antioxidants, and the degenerative diseases of aging. *Proc. Natl. Acad. Sci. USA* 90: 7915-7922.
4. Hayakawa, M., Sugiyama S, Hattori K, Takasawa M, Ozawa T. 1993. Age-associated damage in mitochondrial DNA in human hearts. *Mol. Cell. Biochem.* 119: 95-103.
5. Nishioka, K., Ohtsubo, T., Oda, H., Fujiwara, T., Kang, D., Sugimachi, K. and Nakabeppu, Y. 1999. Expression and differential intracellular localization of two major forms of human 8-oxoguanine DNA glycosylase encoded by alternatively spliced OGG1 mRNAs. *Mol. Biol. Cell* 10: 1637-1652.
6. Boiteux, S. and Radicella, J.P. 2000. The human OGG1 gene: structure, functions, and its implication in the process of carcinogenesis. *Arch. Biochem. Biophys.* 377: 1-8.
7. Minowa, O., Arai, T., Hirano, M., Monden, Y., Nakai, S., Fukuda, M., Itoh, M., Takano, H., Hippou, Y., Aburatani, H., Masumura, K., Nohmi, T., Nishimura, S. and Noda, T. 2000. Mmh/OGG1 gene inactivation results in accumulation of 8-hydroxyguanine in mice. *Proc. Natl. Acad. Sci. USA* 97: 4156-4161.

CHROMOSOMAL LOCATION

Genetic locus: OGG1 (human) mapping to 3p25.3.

PRODUCT

OGG1/2 (h): 293T Lysate represents a lysate of human OGG1/2 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

OGG1/2 (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive OGG1/2 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.