



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

p18 (h): 293T Lysate: sc-174498

BACKGROUND

The normal progression of cells through the cell cycle is under the control of the cyclin dependent protein kinases Cdk4 and Cdk6, which are subject to inhibition by the mitotic inhibitory protein p16. Isolated members of the p16 family have been designated p15 and p18. p15 expression is upregulated approximately 30-fold in TGF β -treated human keratinocytes. The gene encoding p15 has been mapped to chromosome 9p21 at a position adjacent to the p16 gene, at a site of frequent chromosomal abnormality in human tumors. It has been suggested that p15 may function as an effector of TGF β -mediated cell cycle arrest through inhibition of Cdk4 and Cdk6 kinase. The second p16-related protein, p18, interacts strongly with Cdk6 and to a lesser extent with Cdk4, but lacks apparent interaction with other Cdk6s. Recombinant p18 has been shown to inhibit cyclin D-Cdk6 kinase activity. In contrast to p21/p27 that form ternary complexes with cyclin-Cdk6s, only binary complexes of p15, p16 and p18 have been identified in association with Cdk4 and/or Cdk6.

REFERENCES

- Serrano, M., Hannon, G.J. and Beach, D. 1993. A new regulatory motif in cell cycle control causing specific inhibition of cyclin D/CDK4. *Nature* 366: 704-707.
- Sherr, C.J. 1994. G₁ phase progression: cycling on cue. *Cell* 79: 551-555.
- Hunter, T. and Pines, J. 1994. Cyclins and cancer II: cyclin D and CDK inhibitors come of age. *Cell* 79: 573-582.
- Kamb, A., Gruis, N.A., Weaver-Feldhaus, J., Liu, Q., Harshman, K., Tavitgian, S.V., Stockert, E., Day, R.F. III, Johnson, B.E. and Skolnick, M.H. 1994. A cell cycle regulator potentially involved in genesis of many tumor types. *Science* 264: 436-440.
- Hannon, G.J. and Beach, D. 1994. p15INK4B is a potential effector of TGF- β -induced cell cycle arrest. *Nature* 371: 257-261.
- Guan, K.L., Jenkins, C.W., Li, Y., Nichols, M.A., Wu, X., O'Keefe, C.L., Matera, A.G. and Xiong, Y. 1994. Growth suppression by p18, a p16INK4/MTS1 and p14INK4B/MTS2-related CDK6 inhibitor, correlates with wildtype pRb function. *Genes Dev.* 8: 2939-2952.
- Hussussian, C.J., Struewing, J.P., Goldstein, A.M., Higgins, P.A.T., Ally, D.S., Sheahan, M.D., Clark, W.H. Jr., Tucker, M.A. and Dracopoli, N.C. 1994. Germline p16 mutations in familial melanoma. *Nat. Genet.* 8: 15.
- Cairns, P., Mao, L., Merlo, A., Lee, D.J., Schwab, D., Eby, Y., Tokino, K., van der Riet, P., Blaugrund, J.E. and Sidransky, D. 1994. Rates of p16 (MTS1) mutations in primary tumors with 9p loss. *Science* 265: 415-417.

CHROMOSOMAL LOCATION

Genetic locus: CDKN2C (human) mapping to 1p32.3.

PRODUCT

p18 (h): 293T Lysate represents a lysate of human p18 transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

p18 (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive p18 antibodies. Recommended use: 10-20 μ l per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

p18 (DCS-118): sc-56332 is recommended as a positive control antibody for Western Blot analysis of enhanced human p18 expression in p18 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



p18 (DCS-118): sc-56332. Western blot analysis of p18 expression in non-transfected: sc-117752 (A) and human p18 transfected: sc-174498 (B) 293T whole cell lysates.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.