



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# PRP19 (h2): 293T Lysate: sc-174522

## BACKGROUND

The spliceosome, the gigantic molecular machine that performs pre-mRNA splicing in eukaryotes, contains over 200 different proteins and five RNA molecules (U1, U2, U4, U5 and U6). Pre-mRNA splicing is essential to remove internal non-coding regions of pre-mRNA (introns) and to join the remaining segments (exons) into mRNA before translation. The PRP19-associated complex is required for stable association of U5 and U6 with the spliceosome after U4 is released. Changes within the spliceosome upon binding of the PRP19-associated complex include remodeling of the U6/5' splice site interaction and destabilization of Lsm proteins to allow further interaction of U6 with the intron sequence.

## REFERENCES

- Cheng, S.C., Tarn, W.Y., Tsao, T.Y. and Abelson, J. 1993. PRP19: a novel spliceosomal component. *Mol. Cell. Biol.* 13: 1876-1882.
- Tarn, W.Y., Lee, K.R. and Cheng, S.C. 1993. Yeast precursor mRNA processing protein PRP19 associates with the spliceosome concomitant with or just after dissociation of U4 small nuclear RNA. *Proc. Natl. Acad. Sci. USA* 90: 10821-10825.
- Tarn, W.Y., Hsu, C.H., Huang, K.T., Chen, H.R., Kao, H.Y., Lee, K.R. and Cheng, S.C. 1994. Functional association of essential splicing factor(s) with PRP19 in a protein complex. *EMBO J.* 13: 2421-2431.
- Nilsen, T.W. 2003. The spliceosome: the most complex macromolecular machine in the cell? *Bioessays* 25: 1147-1149.
- Turner, I.A., Norman, C.M., Churcher, M.J. and Newman, A.J. 2004. Roles of the U5 snRNP in spliceosome dynamics and catalysis. *Biochem. Soc. Trans.* 32: 928-931.
- Butcher, S.E. and Brow, D.A. 2005. Towards understanding the catalytic core structure of the spliceosome. *Biochem. Soc. Trans.* 33: 447-449.
- Valadkhan, S. 2005. snRNAs as the catalysts of pre-mRNA splicing. *Curr. Opin. Chem. Biol.* 9: 603-608.
- Chan, S.P. and Cheng, S.C. 2005. The Prp19-associated complex is required for specifying and U6 with pre-mRNA during spliceosome activation. *J. Biol. Chem.* 280: 31190-31199.
- Chen, C.H., Kao, D.I., Chan, S.P., Kao, T.C., Lin, J.Y. and Cheng, S.C. 2006. Functional links between the Prp19-associated complex, U4/U6 biogenesis, and spliceosome recycling. *RNA* 12: 765-774.

## CHROMOSOMAL LOCATION

Genetic locus: PRPF19 (human) mapping to 11q12.2.

## PRODUCT

PRP19 (h2): 293T Lysate represents a lysate of human PRP19 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

## STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

## APPLICATIONS

PRP19 (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive PRP19 antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

PRP19 (36-K): sc-101236 is recommended as a positive control antibody for Western Blot analysis of enhanced human PRP19 expression in PRP19 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:  
 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

## DATA



PRP19 (36-K): sc-101236. Western blot analysis of PRP19 expression in non-transfected: sc-117752 (A) and human PRP19 transfected: sc-174522 (B) 293T whole cell lysates.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.