



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien


T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Ribosomal Protein L28 (h2): 293T Lysate: sc-174734

BACKGROUND

The genes encoding for mammalian Ribosomal Proteins comprise multigene families that consist predominantly of multiple processed pseudogenes and one functional intro-containing gene within their coding regions. Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. 60S Ribosomal Protein L28 is encoded by the RPL28 gene. This protein which is a structural constituent of the ribosome, is an RNA binding protein involved in protein biosynthesis.

REFERENCES

1. Gross, T., Nischt, R., Gatermann, K., Swida, U. and Kaufer, N.F. 1988. Primary structure of the ribosomal protein gene S6 from *Schizosaccharomyces pombe*. *Curr. Genet.* 13: 57-63.
2. Lott, J.B. and Mackie, G.A. 1988. Isolation and characterization of cloned cDNAs that code for human Ribosomal Protein S6. *Gene* 65: 31-39.
3. Heinze, H., Arnold, H.H., Fischer, D. and Kruppa, J. 1988. The primary structure of the human Ribosomal Protein S6 derived from a cloned cDNA. *J. Biol. Chem.* 263: 4139-4144.
4. Wool, I.G., Chan, Y.L., Paz, V. and Olvera, J. 1990. The primary structure of rat ribosomal proteins: the amino acid sequences of L27a and L28 and corrections in the sequences of S4 and S12. *Biochim. Biophys. Acta* 1050: 69-73.
5. Feo, S., Davies, B. and Fried, M. 1992. The mapping of seven intron-containing Ribosomal Protein genes shows they are unlinked in the human genome. *Genomics* 13: 201-207.
6. Frigerio, J.M., Dagorn, J.C. and Iovanna, J.L. 1995. Cloning, sequencing and expression of the L5, L21, L27a, L28, S5, S9, S10 and S29 human Ribosomal Protein mRNAs. *Biochim. Biophys. Acta* 1262: 64-68.
7. Hernandez, V.P. and Fallon, A.M. 1999. Ribosomal Protein S6 cDNA from two *Aedes mosquitoes* encodes a carboxyl-terminal extension that resembles Histone H1 proteins. *Genetica* 106: 263-267.

CHROMOSOMAL LOCATION

Genetic locus: RPL28 (human) mapping to 19q13.42.

PRODUCT

Ribosomal Protein L28 (h2): 293T Lysate represents a lysate of human Ribosomal Protein L28 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

APPLICATIONS

Ribosomal Protein L28 (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive Ribosomal Protein L28 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.