



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



# CD33 (h4): 293T Lysate: sc-174855

## BACKGROUND

CD33 is a type I transmembrane glycoprotein that is found on granulocyte and macrophage precursors in the bone marrow, but is absent from pluripotent stem cells. CD33 is also expressed on monocytes in peripheral blood. It is used as a marker to distinguish myelogenous leukemia cells from lymphoid or erythroid leukemias. CD33 may function as a sialic acid-dependent cell adhesion molecule.

## REFERENCES

- Griffin, J.D., Linch, D., Sabbath, K., Larcom, P. and Schlossman, S.F. 1984. A monoclonal antibody reactive with normal and leukemic human myeloid progenitor cells. *Leuk. Res.* 8: 521-534.
- Favaloro, E.J., Bradstock, K.F., Kabral, A., Grimsley, P. and Berndt, M.C. 1987. Characterization of monoclonal antibodies to the human myeloid-differentiation antigen, "gp67" (CD33). *Dis. Markers* 5: 215-225.
- Andrews, R.G., Singer, J.W. and Bernstein, I.D. 1989. Precursors of colony-forming cells in humans can be distinguished from colony-forming cells by expression of the CD33 and CD34 antigens and light scatter properties. *J. Exp. Med.* 169: 1721-1731.
- Handgretinger, R., Schafer, H.J., Baur, F., Frank, D., Ottenlinger, C., Buhning, H.J. and Niethammer, D. 1993. Expression of an early myelopoietic antigen (CD33) of a subset of human umbilical cord blood-derived natural killer cells. *Immunol. Lett.* 37: 223-228.
- Pierelli, L., Teofili, L., Menichella, G., Rumi, C., Paolini, A., Iovino, S., Puggioni, P.L., Leone, G. and Bizzi, B. 1993. Further investigations on the expression of HLA-DR, CD33 and CD13 surface antigens in purified bone marrow and peripheral blood CD34<sup>+</sup> haematopoietic progenitor cells. *Br. J. Haematol.* 84: 24-30.
- Freeman, S.D., Kelm, S., Barber, E.K. and Crocker, P.R. 1995. Characterization of CD33 as a new member of the sialoadhesin family of cellular interaction molecules. *Blood* 85: 2005-2012.
- Kelm, S., Schauer, R. and Crocker, P.R. 1996. The Sialoadhesins: a family of sialic-acid-dependent cellular recognition molecules within the immunoglobulin superfamily. *Glycoconj. J.* 13: 913-926.
- Sgroi, D., Nocks, A. and Stamenkovic, I. 1996. A single N-linked glycosylation site is implicated in the regulation of ligand recognition by the I-type lectins CD22 and CD33. *J. Biol. Chem.* 271: 18803-18809.
- Vitale, C., Romagnani, C., Puccetti, A., Olive, D., Costello, R., Chiossone, L., Pitto, A., Bacigalupo, A., Moretta, L. and Mingari, M.C. 2001. Surface expression and function of p75/AIRM-1 or CD33 in acute myeloid leukemias: engagement of CD33 induces apoptosis of leukemic cells. *Proc. Natl. Acad. Sci. USA* 98: 5764-5769.

## CHROMOSOMAL LOCATION

Genetic locus: CD33 (human) mapping to 19q13.41.

## PRODUCT

CD33 (h4): 293T Lysate represents a lysate of human CD33 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

## APPLICATIONS

CD33 (h4): 293T Lysate is suitable as a Western Blotting positive control for human reactive CD33 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

## STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.