



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

CIN85 (h2): 293T Lysate: sc-175351

BACKGROUND

CD2AP (CMS) and CIN85 (Ruk) belong to a family of ubiquitously expressed adaptor molecules containing three SH3 domains, a proline-rich region and a coiled-coil domain. By binding to numerous proteins, CD2AP and CIN85 assemble multimeric complexes implicated in cell-specific signals controlling T-cell activation, kidney glomeruli function or apoptosis in neuronal cells. CIN85/CD2AP also associate with accessory endocytic proteins, components of the actin cytoskeleton, and other adaptor proteins involved in receptor tyrosine kinase signaling. These interactions enable CIN85/CD2AP to function within a network of signaling pathways that coordinate critical steps involved in downregulation and degradation of receptor tyrosine kinases.

REFERENCES

1. Take, H., et al. 2000. Cloning and characterization of a novel adaptor protein, CIN85, that interacts with c-Cbl. *Biochem. Biophys. Res. Commun.* 268: 321-328.
2. Watanabe, S., et al. 2000. Characterization of the CIN85 adaptor protein and identification of components involved in CIN85 complexes. *Biochem. Biophys. Res. Commun.* 278: 167-174.
3. Szymkiewicz, I., et al. 2002. CIN85 participates in Cbl- β -mediated down-regulation of receptor tyrosine kinases. *J. Biol. Chem.* 277: 39666-39672.
4. Haglund, K., et al. 2002. Cbl-directed monoubiquitination of CIN85 is involved in regulation of ligand-induced degradation of EGF receptors. *Proc. Natl. Acad. Sci. USA* 99: 12191-12196.
5. Dikic, I. 2002. CIN85/CMS family of adaptor molecules. *FEBS Lett.* 529: 110-115.
6. Schmidt, M.H., et al. 2003. SETA/CIN85/Ruk and its binding partner AIP1 associate with diverse cytoskeletal elements, including FAKs and modulate cell adhesion. *J. Cell Sci.* 116: 2845-2855.

CHROMOSOMAL LOCATION

Genetic locus: SH3KBP1 (human) mapping to Xp22.12.

PRODUCT

CIN85 (h2): 293T Lysate represents a lysate of human CIN85 transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

CIN85 (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive CIN85 antibodies. Recommended use: 10-20 μ l per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

CIN85 (C-8): sc-271567 is recommended as a positive control antibody for Western Blot analysis of enhanced human CIN85 expression in CIN85 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



CIN85 (C-8): sc-271567. Western blot analysis of CIN85 expression in non-transfected: sc-117752 (A) and human CIN85 transfected: sc-175351 (B) 293T whole cell lysates.



CIN85 (G-12): sc-271310. Western blot analysis of CIN85 expression in non-transfected: sc-117752 (A) and human CIN85 transfected: sc-175351 (B) 293T whole cell lysates.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.