



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

S-100A15 (h3): 293T Lysate: sc-175804

BACKGROUND

Psoriasin, also known as PSOR1 or S-100A7, is a 101 amino acid protein that belongs to the S-100 family of calcium binding proteins and is secreted via a non-classical secretory pathway into the cytoplasm. Expressed in fetal ear, tongue and skin, Psoriasin is thought to function in the regulation of many cellular processes, including the cell cycle, cell progression and cellular differentiation. Psoriasin contains two EF-hand domains and is highly upregulated in psoriatic epidermis, as well as in bladder squamous cell carcinoma and breast cancer tissue, suggesting a possible role in carcinogenesis. The gene encoding Psoriasin and the related S100A15 gene are thought to have diverged from one mouse gene, designated S100A15. In humans, the S100A15 gene encodes a calcium binding protein, also known as S-100A7A, that shares 95% sequence identity with Psoriasin.

REFERENCES

1. Brodersen, D.E., et al. 1998. EF-hands at atomic resolution: the structure of human psoriasin (S100A7) solved by MAD phasing. *Structure* 6: 477-489.
2. Ruse, M., et al. 2003. S100A7 (psoriasin) interacts with epidermal fatty acid binding protein and localizes in focal adhesion-like structures in cultured keratinocytes. *J. Invest. Dermatol.* 121: 132-141.
3. Wolf, R., et al. 2003. Molecular cloning and characterization of alternatively spliced mRNA isoforms from psoriatic skin encoding a novel member of the S100 family. *FASEB J.* 17: 1969-1971.
4. Jiang, W.G., et al. 2004. Psoriasin is aberrantly expressed in human breast cancer and is related to clinical outcomes. *Int. J. Oncol.* 25: 81-85.
5. Martinsson, H., et al. 2005. Expression patterns of S100A7 (psoriasin) and S100A9 (calgranulin-B) in keratinocyte differentiation. *Exp. Dermatol.* 14: 161-168.
6. Webb, M., et al. 2005. Expression analysis of the mouse S100A7/psoriasin gene in skin inflammation and mammary tumorigenesis. *BMC Cancer* 5: 17.
7. Porre, S., et al. 2005. Psoriasin, a calcium-binding protein with chemotactic properties is present in the third trimester amniotic fluid. *Mol. Hum. Reprod.* 11: 87-92.
8. Eckert, R.L., et al. 2006. S100A7 (Psoriasin): a story of mice and men. *J. Invest. Dermatol.* 126: 1442-1444.
9. Wolf, R., et al. 2006. The mouse S100A15 ortholog parallels genomic organization, structure, gene expression, and protein-processing pattern of the human S100A7/A15 subfamily during epidermal maturation. *J. Invest. Dermatol.* 126: 1600-1608.

CHROMOSOMAL LOCATION

Genetic locus: S100A7 (human) mapping to 1q21.3.

PRODUCT

S-100A15 (h3): 293T Lysate represents a lysate of human S-100A15 transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

S-100A15 (h3): 293T Lysate is suitable as a Western Blotting positive control for human reactive S-100A15 antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.