



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

IL-21R (h2): 293T Lysate: sc-175935

BACKGROUND

The IL-21 receptor (also designated IL-21R, NILR or novel interleukin receptor) is a type I cytokine receptor that forms a complex with the cytokine receptor γ chain, $\gamma(c)$, and mediates IL-21 signaling. IL-21R is present on the surface of natural killer, B and T cell populations with high levels in spleen and thymus. IL-21 and IL-21R influence lymphoid proliferation and early lymphoid development in the transition between innate and adaptive immunity. Tumor necrosis factor (TNF) upregulates IL-21R, and combinations of TNF and IL-21 can have synergistic effects on myeloma cell proliferation through pathways involving phosphorylation of JAK1, Stat3 and ERK 1/2. The human IL-21R gene maps to chromosome 16p11 and encodes a 538 amino acid protein that is closely related to human IL2RB and shares 62% sequence identity to mouse IL21r.

REFERENCES

1. Parrish-Novak, J., et al. 2000. Interleukin-21 and its receptor are involved in NK cell expansion and regulation of lymphocyte function. *Nature* 408: 57-63.
2. Asao, H., et al. 2001. Cutting edge: the common γ chain is an indispensable subunit of the IL-21 receptor complex. *J. Immunol.* 167: 1-5.
3. Vosshenrich, C.A. and Di Santo, J.P. 2001. Cytokines: IL-21 joins the γ_c -dependent network? *Curr. Biol.* 11: R175-R177.
4. Habib, T., et al. 2002. The common γ chain (γ_c) is a required signaling component of the IL-21 receptor and supports IL-21-induced cell proliferation via JAK3. *Biochemistry* 41: 8725-8731.
5. Kasaian, M.T., et al. 2002. IL-21 limits NK cell responses and promotes antigen-specific T cell activation: a mediator of the transition from innate to adaptive immunity. *Immunity* 16: 559-569.
6. Ueda, C., et al. 2002. The gene for interleukin-21 receptor is the partner of BCL6 in t(3;16)(q27;p11), which is recurrently observed in diffuse large B cell lymphoma. *Oncogene* 21: 368-376.
7. Brenne, A.T., et al. 2002. Interleukin-21 is a growth and survival factor for human myeloma cells. *Blood* 99: 3756-3762.
8. Online Mendelian Inheritance in Man, OMIM[™]. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 605383. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
9. LocusLink Report (LocusID: 50615). <http://www.ncbi.nlm.nih.gov/LocusLink/>

CHROMOSOMAL LOCATION

Genetic locus: IL21R (human) mapping to 16p12.1.

PRODUCT

IL-21R (h2): 293T Lysate represents a lysate of human IL-21R transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

IL-21R (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive IL-21R antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.