



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

GFR α -3 (h2): 293T Lysate: sc-176275

BACKGROUND

Glial cell line-derived neurotrophic factor (GDNF) and the related neurotrophic factor neurturin (NTN) are potent survival factors for central and peripheral neurons. GDNF is a glycosylated, disulfide-bonded homodimer that is distantly related to the TGF β superfamily of growth factors. Three receptors for these factors, GFR α -1 (also designated GDNFR- α , RETL1 or TrnR-1), GFR α -2 (also designated GDNFR- β , RETL2, NTN- α or TrnR-2) and GFR α -3 have been identified. The receptors do not contain transmembrane domains and are attached to the cell membrane by glycosyl-phosphoinositol linkage. Both GFR α -1 and GFR α -2 have been shown to mediate the GDNF-dependent and NTN-dependent phosphorylation and activation of the tyrosine kinase Ret. GFR α -3 is expressed only during development.

REFERENCES

1. Lin, L.F., Doherty, D.H., Lile, J.D., Bektesh, S. and Collins, F. 1993. GDNF: a glial cell line-derived neurotrophic factor for midbrain dopaminergic neurons. *Science* 260: 1130-1132.
2. Jing, S., Wen, D., Yu, Y., Holst, P.L., Luo, Y., Fang, M., Tamir, R., Antonio, L., Hu, Z., Cupples, R., Louis, J.C., Hu, S., Altmann, B.W. and Fox, G.M. 1996. GDNF-induced activation of the Ret protein tyrosine kinase is mediated by GDNFR- α , a novel receptor for GDNF. *Cell* 85: 1113-1124.
3. Treanor, J.J., Goodman, L., de Sauvage, F., Stone, D.M., Poulsen, K.T., Beck, C.D., Gray, C., Armanini, M.P., Pollock, R.A., Hefti, F., Phillips, H.S., Goddard, A., Moore, M.W., Buj-Bello, A., Davies, A.M., et al. 1996. Characterization of a multi-component receptor for GDNF. *Nature* 382: 80-83.
4. Kotzbauer, P.T., Lampe, P.A., Heuckeroth, R.O., Golden, J.P., Creedon, D.J., Johnson, E.M., Jr. and Milbrandt, J. 1996. Neurturin, a relative of glial-cell-line-derived neurotrophic factor. *Nature* 384: 467-470.
5. Baloh, R.H., Tansey, M.G., Golden, J.P., Creedon, D.J., Heuckeroth, R.O., Keck, C.L., Zimonjic, D.B., Popescu, N.C., Johnson, E.M., Jr. and Milbrandt, J. 1997. TrnR-2, a novel receptor that mediates neurturin and GDNF signaling through Ret. *Neuron* 18: 793-802.
6. Naveilhan, P., Baudet, C., Mikaelis, A., Shen, L., Westphal, H. and Ernfors, P. 1998. Expression and regulation of GFR α -3, a glial cell line-derived neurotrophic factor family receptor. *Proc. Natl. Acad. Sci. USA* 95: 1295-1300.

CHROMOSOMAL LOCATION

Genetic locus: GFRA3 (human) mapping to 5q31.2.

PRODUCT

GFR α -3 (h): 293T Lysate represents a lysate of human GFR α -3 transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

STORAGE

Store at -20 $^{\circ}$ C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

GFR α -3 (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive GFR α -3 antibodies.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.