



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

c-Src (h2): 293T Lysate: sc-176937

BACKGROUND

The major translational products of the Src gene family are membrane-associated tyrosine protein kinases that lack transmembrane and external amino acid sequences. By virtue of their common structural motifs, the Src family is composed of nine members in vertebrates, including c-Src, c-Yes, c-Fgr, Yrk, Fyn, Lyn, Hck, Lck and Blk. Src family kinases, which contain an amino-terminal cell membrane anchor followed by SH3 and SH2 domains, transduce signals that are involved in the control of a variety of cellular processes, including proliferation, differentiation, motility and adhesion. Src family members are normally maintained in an inactive state and can be activated transiently during cellular events such as mitosis. Different subcellular locations of Src family kinases may be important for the regulation of specific cellular processes, such as mitogenesis, cytoskeletal organization and membrane trafficking. c-Src (also designated pp60Src, Src p60 and proto-oncogene tyrosine protein kinase Src) is expressed in a broad range of tissue and cell types, although the highest levels of c-Src are detected in neuronal tissues and platelets. c-Src may play a role in events associated with both neuronal differentiation and maintenance of mature neuronal cell functions.

REFERENCES

1. Sakaguchi, A.Y. 1983. Organization of human proto-oncogenes. *Prog. Clin. Biol. Res.* 119: 93-103.
2. Brugge, J.S., Cotton, P.C., Quesada, A.E., Barret, J.N., Nonner, D. and Keane, R.W. 1985. Neurons express high levels of structurally modified, activated form of pp60Src. *Nature* 316: 554-557.
3. Golden, A., Nemeth, S.P. and Brugge, J.S. 1986. Bloodplatelets express high levels of the pp60Src-specific tyrosine kinase activity. *Proc. Natl. Acad. Sci. USA* 83: 852-856.
4. Cartwright, C.A., Simantov, R., Kaplan, P.L., Hunter, T. and Eckhart, W. 1987. Alterations in pp60Src accompany differentiation of neurons from rat embryo striatum. *Mol. Cell. Biol.* 7: 1830-1840.
5. Wiestler, O.D. and Walter, G. 1988. Developmental expression of two forms of pp60Src in mouse brain. *Mol. Cell. Biol.* 8: 502-504.
6. Eiseman, E. and Bolen, J.B. 1990. Src-related tyrosine protein kinases as signaling components in hematopoietic cells. *Cancer Cells* 2: 303-310.
7. Bolen, J.B., Thompson, P.A., Eiseman, E. and Horak, I.D. 1991. Expression and interactions of the Src family of tyrosine protein kinases in T lymphocytes. *Adv. Cancer Res.* 57: 103-149.
8. Broome, M.A. and Hunter, T. 1997. The PDGF receptor phosphorylates Tyr 138 in the c-Src domain *in vivo* reducing peptide ligand binding. *Oncogene* 14: 17-34.
9. Gilmore, E.S., Stutts, M.J. and Milgram, S.L. 2001. Src family kinases mediate epithelial Na⁺ channel inhibition by endothelin. *J. Biol. Chem.* 276: 42610-42617.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

CHROMOSOMAL LOCATION

Genetic locus: SRC (human) mapping to 20q11.23.

PRODUCT

c-Src (h2): 293T Lysate represents a lysate of human c-Src transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.


APPLICATIONS

c-Src (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive c-Src antibodies. Recommended use: 10-20 µl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

c-Src (17AT28): sc-130124 is recommended as a positive control antibody for Western Blot analysis of enhanced human c-Src expression in c-Src transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

DATA



c-Src (17AT28): sc-130124. Western blot analysis of c-Src expression in non-transfected: sc-117752 (A) and human c-Src transfected: sc-176937 (B) 293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.