

# Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

#### SANTA CRUZ BIOTECHNOLOGY, INC.

## Fos B (h2): 293T Lysate: sc-177246



#### BACKGROUND

The v-Fos oncogene was initially identified as the transforming gene of two independent murine osteosarcoma virus isolates and an avian nephroblastoma virus. The cellular homolog, c-Fos, encodes a nuclear phosphoprotein that is rapidly and transiently induced by a variety of agents and functions as a transcriptional regulator for several genes. In contrast to c-Jun proteins, which form homo- and heterodimers which bind to specific DNA TPA response elements (TREs), c-Fos proteins are only active as heterodimers with members of the Jun gene family. Murine Fos B encodes a nuclear protein of 338 amino acids which has 70% homology with c-Fos, exhibits similar kinetics of expression as c-Fos and forms heterodimers with both c-Jun and Jun B, which bind to TRE DNA response elements. Functional homologs of c-Fos and Fos B include Fra-1 and Fra-2 genes.

#### REFERENCES

- Finkel, M.P., et al. 1966. Virus induction of osteosarcomas in mice. Science 151: 698-701.
- Curran, T., et al. 1984. FBR murine osteosarcoma virus. I. Molecular analysis and characterization of a 75,000-Da gag-fos fusion product. Virology 135: 218-228.
- Sambucetti, L.C., et al. 1986. The fos protein complex is associated with DNA in isolated nuclei and binds to DNA cellulose. Science 234: 1417-1419.
- Nishizawa, M., et al. 1987. An avian transforming retrovirus isolated from a nephroblastoma that carries the fos gene as the oncogene. J. Virol. 61: 3733-3740.
- Renz, M., et al. 1987. Chromatin association and DNA-binding properties of the c-Fos protooncogene product. Nucleic Acids Res. 15: 277-292.
- Zerial, M., et al. 1989. The product of a novel growth factor activated gene, Fos B, interacts with JUN proteins enhancing their DNA binding activity. EMBO J. 8: 805-813.
- Nishina, H., et al. 1990. Isolation and characterization of Fra-2, an additional member of the fos gene family. Proc. Natl. Acad. Sci. USA 87: 3619-3623.
- 8. Castellazzi, M., et al. 1991. Overexpression of c-Jun, Jun B, or Jun D affects cell growth differently. Proc. Natl. Acad. Sci. USA 88: 8890-8894.

#### CHROMOSOMAL LOCATION

Genetic locus: FOSB (human) mapping to 19q13.32.

#### PRODUCT

Fos B (h2): 293T Lysate represents a lysate of human Fos B transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

#### **STORAGE**

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

#### **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

#### APPLICATIONS

Fos B (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive Fos B antibodies. Recommended use:  $10-20 \mu$ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

Fos B (B-10): sc-390035 is recommended as a positive control antibody for Western Blot analysis of enhanced human Fos B expression in Fos B transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

#### **RECOMMENDED SUPPORT REAGENTS**

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker<sup>TM</sup> Molecular Weight Standards: sc-2035, UltraCruz<sup>®</sup> Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

#### DATA





Fos B (B-10): sc-390035. Western blot analysis of Fos B expression in non-transfected: sc-117752 (**A**) and human Fos B transfected: sc-177246 (**B**) 293T whole cell lysates. Fos B (83B1138): sc-52926. Western blot analysis of Fos B expression in non-transfected: sc-117752 (A) and human Fos B transfected: sc-177246 (B) 293T whole cell lysates.

#### **PROTOCOLS**

See our web site at www.scbt.com for detailed protocols and support products.