



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

LRRFIP1 (h3): 293T Lysate: sc-177274

BACKGROUND

LRRFIP1 (also designated GCF2) is a 738 amino acid human protein whose rodent counterpart is known as *Lrrfip1* (also designated FLAP in mouse). LRRFIP1 is a transcriptional repressor which will preferentially bind to the GC-rich consensus sequence (5'-AGCCCCGGCG-3') and may also regulate expression of TNF, EGFR and PDGF-A. LRRFIP1 is also believed to control smooth muscle cell proliferation following arterial injury through PDGF-A repression. The N-terminus of LRRFIP1 shows high homology to the coiled-coil domain of FLAP, a protein which binds the leucine-rich repeat (LRR) of Flightless I, and the interaction of LRRFIP1 with the LRR of Flightless I has been confirmed. LRRFIP1 does not bind single-stranded DNA or RNA significantly and binds double-stranded DNA weakly. In contrast, LRRFIP1 binds double-stranded RNA with high affinity, and two molecules of LRRFIP1 bind the TaR stem. The RNA binding domain has been identified and encompasses a lysine-rich motif. Flightless I has a C-terminal TaR-like domain which binds Actin and therefore the association of LRRFIP1 with the LRR of Flightless I may provide a link between the Actin cytoskeleton and RNA in mammalian cells.

REFERENCES

1. Reed, A.L., Yamazaki, H., Kaufman, J.D., Rubinstein, Y., Murphy, B. and Johnson, A.C. 1998. Molecular cloning and characterization of a transcription regulator with homology to GC-binding factor. *J. Biol. Chem.* 273: 21594-21602.
2. Wilson, S.A., Brown, E.C., Kingsman, A.J. and Kingsman, S.M. 1998. TRIP: a novel double stranded RNA binding protein which interacts with the leucine rich repeat of Flightless I. *Nucleic Acids Res.* 26: 3460-3467.
3. Khachigian, L.M., Santiago, F.S., Rafty, L.A., Chan, O.L., Delbridge, G.J., Bobik, A., Collins, T. and Johnson, A.C. 1999. GC factor 2 represses platelet-derived growth factor A-chain gene transcription and is itself induced by arterial injury. *Circ. Res.* 84: 1258-1267.
4. Rikiyama, T., Curtis, J., Oikawa, M., Zimonjic, D.B., Popescu, N., Murphy, B.A., Wilson, M.A. and Johnson, A.C. 2003. GCF2: expression and molecular analysis of repression. *Biochim. Biophys. Acta* 1629: 15-25.
5. Suriano, A.R., Sanford, A.N., Kim, N., Oh, M., Kennedy, S., Henderson, M.J., Dietzmann, K. and Sullivan, K.E. 2005. GCF2/LRRFIP1 represses tumor necrosis factor α expression. *Mol. Cell. Biol.* 25: 9073-9081.
6. Sjöblom, T., Jones, S., Wood, L.D., Parsons, D.W., Lin, J., Barber, T.D., Mandelker, D., Leary, R.J., Ptak, J., Silliman, N., Szabo, S., Buckhaults, P., Farrell, C., Meeh, P., Markowitz, S.D., Willis, J., Dawson, D., Willson, J.K., Gazdar, A.F., Hartigan, J., Wu, L., Liu, C., Parmigiani, G., Park, B.H., et al. 2006. The consensus coding sequences of human breast and colorectal cancers. *Science* 314: 268-274.

CHROMOSOMAL LOCATION

Genetic locus: LRRFIP1 (human) mapping to 2q37.3.

PRODUCT

LRRFIP1 (h3): 293T Lysate represents a lysate of human LRRFIP1 transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

LRRFIP1 (h3): 293T Lysate is suitable as a Western Blotting positive control for human reactive LRRFIP1 antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.