

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PDEF (h2): 293T Lysate: sc-177713



The Power to Question

BACKGROUND

Prostate epithelium-specific Ets transcription factor (PDEF), also designated prostate Ets or SAM pointed domain-containing Ets transcription factor, is a 335 amino acid nuclear protein. PDEF belongs to the Ets family of proteins. This protein, which localizes to prostate epithelial cells, functions as an Ets transcription factor. It upregulates the activity of the p62 promoter but this activity can be downregulated by PSI. It is also involved in the activation of prostate-specific antigen (PSA) by acting as an androgen-independent transactivator.

REFERENCES

- Oettgen, P., et al. 2000. PDEF, a novel prostate epithelium-specific Ets transcription factor, interacts with the androgen receptor and activates prostate-specific antigen gene expression. J. Biol. Chem. 275: 1216-1225.
- Yamada, N., et al. 2000. Cloning and expression of the mouse Pse gene encoding a novel Ets family member. Gene 241: 267-274.
- Chen, H., et al. 2002. Nkx-3.1 interacts with prostate-derived Ets factor and regulates the activity of the PSA promoter. Cancer Res. 62: 338-340.
- Feldman, R.J., et al. 2003. PDEF expression in human breast cancer is correlated with invasive potential and altered gene expression. Cancer Res. 63: 4626-4631.
- 5. Thompson, H.G., et al. 2003. p62 overexpression in breast tumors and regulation by prostate-derived Ets factor in breast cancer cells. Oncogene 22: 2322-2333.
- Chen, H., et al. 2005. Structural and functional analysis of domains mediating interaction between Nkx-3.1 and PDEF. J. Cell. Biochem. 94: 168-177.
- 7. Wang, Y., et al. 2005. Analysis of the 2.0A crystal structure of the protein-DNA complex of the human PDEF Ets domain bound to the prostate specific antigen regulatory site. Biochemistry 44: 7095-7106.

CHROMOSOMAL LOCATION

Genetic locus: SPDEF (human) mapping to 6p21.31.

PRODUCT

PDEF (h2): 293T Lysate represents a lysate of human PDEF transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

PDEF (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive PDEF antibodies. Recommended use: 10-20 µl per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

PDEF (G-10): sc-166846 is recommended as a positive control antibody for Western Blot analysis of enhanced human PDEF expression in PDEF transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-lgG κ BP-HRP: sc-516102 or m-lgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz MarkerTM Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



PDEF (G-10): sc-166846. Western blot analysis of PDEF expression in non-transfected: sc-117752 (A) and human PDEF transfected: sc-177713 (B) 293T whole cell lysates

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3800 fax 831.457.3801 Europe +00800 4573 8000 49 6221 4503 0 www.scbt.com