



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Suppressin (h): 293T Lysate: sc-178000

BACKGROUND

Suppressin, also known as DEAF-1 (deformed epidermal autoregulatory factor-1), SPN, NUDR (nuclear DEAF-1-related transcriptional regulator), or ZMYND5 (zinc finger MYND domain-containing protein 5), is a transcription factor required for embryonic development. Suppressin contains one SAND domain and one C-terminal MYND-type zinc finger. It interacts with LMO4 and CLIM-2, suggesting that it plays a role mediating cell fate and embryonic pattern formation. Suppressin is expressed in a variety of tissues and localizes to the nucleus. Several isoforms exist due to alternative splicing and, depending on the isoform, Suppressin is secreted in some cell types. Secreted Suppressin can function to inhibit cell proliferation, arresting cells in the G₀ or G₁ phase. Mutations in the gene encoding Suppressin may result in a growth advantage leading to the development and progression of neoplasia. This suggests that Suppressin is a potential target for cancer therapy.

REFERENCES

- Huggenvik, J.I., Michelson, R.J., Collard, M.W., Ziemba, A.J., Gurley, P. and Mowen, K.A. 1998. Characterization of a nuclear deformed epidermal autoregulatory factor-1 (DEAF-1)-related (NUDR) transcriptional regulator protein. *Mol. Endocrinol.* 12: 1619-1639.
- Michelson, R.J., Collard, M.W., Ziemba, A.J., Persinger, J., Bartholomew, B. and Huggenvik, J.I. 1999. Nuclear DEAF-1-related (NUDR) protein contains a novel DNA binding domain and represses transcription of the heterogeneous nuclear ribonucleoprotein A2/B1 promoter. *J. Biol. Chem.* 274: 30510-30519.
- Bottomley, M.J., Collard, M.W., Huggenvik, J.I., Liu, Z., Gibson, T.J. and Sattler, M. 2001. The SAND domain structure defines a novel DNA-binding fold in transcriptional regulation. *Nat. Struct. Biol.* 8: 626-633.
- Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 602635. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Lemondé, S., Turecki, G., Bakish, D., Du, L., Hrdina, P.D., Bown, C.D., Sequeira, A., Kushwaha, N., Morris, S.J., Basak, A., Ou, X.M. and Albert, P.R. 2003. Impaired repression at a 5-hydroxytryptamine 1A receptor gene polymorphism associated with major depression and suicide. *J. Neurosci.* 23: 8788-8799.
- Jensik, P.J., Huggenvik, J.I. and Collard, M.W. 2004. Identification of a nuclear export signal and protein interaction domains in deformed epidermal autoregulatory factor-1 (DEAF-1). *J. Biol. Chem.* 279: 32692-32699.
- Choi, K.O., Lee, T., Lee, N., Kim, J.H., Yang, E.G., Yoon, J.M., Kim, J.H., Lee, T.G. and Park, H. 2005. Inhibition of the catalytic activity of hypoxia-inducible factor-1 α -prolyl-hydroxylase 2 by a MYND-type zinc finger. *Mol. Pharmacol.* 68: 1803-1809.
- Czesak, M., Lemondé, S., Peterson, E.A., Rogaeva, A. and Albert, P.R. 2006. Cell-specific repressor or enhancer activities of DEAF-1 at a Serotonin 1A receptor gene polymorphism. *J. Neurosci.* 26: 1864-1871.
- Spadaccini, R., Perrin, H., Bottomley, M.J., Ansieau, S. and Sattler, M. 2006. Structure and functional analysis of the MYND domain. *J. Mol. Biol.* 358: 498-508.

CHROMOSOMAL LOCATION

Genetic locus: DEAF1 (human) mapping to 11p15.5.

PRODUCT

Suppressin (h): 293T Lysate represents a lysate of human Suppressin transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

Suppressin (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive Suppressin antibodies. Recommended use: 10-20 μ l per lane.


Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

Suppressin (C-8): sc-398387 is recommended as a positive control antibody for Western Blot analysis of enhanced human Suppressin expression in Suppressin transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



Suppressin (C-8): sc-398387. Western blot analysis of Suppressin expression in non-transfected: sc-117752 (A) and human Suppressin transfected: sc-178000 (B) 293T whole cell lysates.



Suppressin (E-5): sc-514180. Western blot analysis of Suppressin expression in non-transfected: sc-117752 (A) and human Suppressin transfected: sc-178000 (B) 293T whole cell lysates.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.