



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TH-POK (h6): 293T Lysate: sc-178046

BACKGROUND

TH-POK (T-helper-inducing POZ/Krüppel-like factor), also known as zinc finger protein 67 (ZFP67), zinc finger and BTB domain-containing protein 7B or Krüppel-related zinc finger protein cKrox, is a 539 amino acid protein that contains one BTB (POZ) domain and four C₂H₂-type zinc fingers. Localized to the nucleus, TH-POK functions primarily as a key regulator of lineage commitment of immature T cell precursors. Specifically, the presence of TH-POK directs positively selected thymocytes to the CD4 lineage, whereas its absence causes default development to the CD8 lineage. TH-POK also functions as a transcriptional repressor of various other genes, such as COL1A1, COL1A2 and Fibronectin.

REFERENCES

1. He, X., He, X., Dave, V.P., Zhang, Y., Hua, X., Nicolas, E., Xu, W., Roe, B.A. and Kappes, D.J. 2005. The zinc finger transcription factor Th-POK regulates CD4 versus CD8 T-cell lineage commitment. *Nature* 433: 826-833.
2. He, X. and Kappes, D.J. 2006. CD4/CD8 lineage commitment: light at the end of the tunnel? *Curr. Opin. Immunol.* 18: 135-142.
3. Kappes, D.J., He, X. and He, X. 2006. Role of the transcription factor Th-POK in CD4:CD8 lineage commitment. *Immunol. Rev.* 209: 237-252.
4. Kimura, H., Morii, E., Ikeda, J.I., Ezoe, S., Xu, J.X., Nakamichi, N., Tomita, Y., Shibayama, H., Kanakura, Y. and Aozasa, K. 2006. Role of DNA methylation for expression of novel stem cell marker CD31 in hematopoietic cells. *Leukemia* 20: 1551-1556.
5. He, X., Park, K., Wang, H., He, X., Zhang, Y., Hua, X., Li, Y. and Kappes, D.J. 2008. CD4-CD8 lineage commitment is regulated by a silencer element at the ThPOK transcription-factor locus. *Immunity* 28: 346-358.
6. Bell, J.J. and Bhandoola, A. 2008. Putting ThPOK in place. *Nat. Immunol.* 9: 1095-1097.
7. Wang, L., Wildt, K.F., Zhu, J., Zhang, X., Feigenbaum, L., Tessarollo, L., Paul, W.E., Fowlkes, B.J. and Bosselut, R. 2008. Distinct functions for the transcription factors GATA-3 and ThPOK during intrathymic differentiation of CD4⁺ T cells. *Nat. Immunol.* 9: 1122-1130.
8. Egawa, T. and Littman, D.R. 2008. ThPOK acts late in specification of the helper T cell lineage and suppresses Runx-mediated commitment to the cytotoxic T cell lineage. *Nat. Immunol.* 9: 1131-1139.
9. Online Mendelian Inheritance in Man, OMIM™. 2008. Johns Hopkins University, Baltimore, MD. MIM Number: 607646. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: ZBTB7B (human) mapping to 1q21.3.

PRODUCT

TH-POK (h6): 293T Lysate represents a lysate of human TH-POK transfected 293T cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

TH-POK (h6): 293T Lysate is suitable as a Western Blotting positive control for human reactive TH-POK antibodies.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.